

From the DEPARTMENT OF ONCOLOGY-PATHOLOGY  
Karolinska Institutet, Stockholm, Sweden

**BETA-ARRESTINS IN CANCER: LINKING PRO-  
TUMORIGENIC EXTRACELLULAR ACTIVATED  
SIGNALING WITH THE TUMOR SUPPRESSOR P53  
PATHWAY**

Naida Suleymanova



**Karolinska  
Institutet**

Stockholm 2017

Cover: *Mosaic  $\beta$ -arrestin isoforms*. Generated through PDB and GIMP.

All previously published papers were reproduced with permission from the publisher.

Published by Karolinska Institutet.

Printed by E-Print AB, 2017

© Naida Suleymanova, 2017

ISBN 978-91-7676-731-3

# **BETA-ARRESTINS IN CANCER: LINKING PRO-TUMORIGENIC EXTRACELLULAR ACTIVATED SIGNALING WITH THE TUMOR SUPPRESSOR P53 PATHWAY**

THESIS FOR DOCTORAL DEGREE (Ph.D.)

By

**Naida Suleymanova, M.D.**

*Principal Supervisor:*

Leonard Girnita, Associate Professor  
Karolinska Institutet  
Department of Oncology-Pathology

*Co-supervisor:*

Klas Wiman, Professor  
Karolinska Institutet  
Department of Oncology-Pathology

*Opponent:*

Gunnar Schulte, Professor  
Karolinska Institutet  
Department of Physiology and Pharmacology

*Examination Board:*

Sergiu-Bogdan Catrina, Associate Professor  
Karolinska Institutet  
Department of Molecular Medicine and Surgery

Cecilia Williams, Professor  
KTH – Royal Institute of Technology  
School of Biotechnology

Anna-Karin Olsson, Associate Professor  
Uppsala University  
Department of Medical Biochemistry and Microbiology



Asya M



## ABSTRACT

The IGF-1R is an important player in cancer development that maintains the malignant phenotype by inducing cell proliferation, survival, transformation, motility and invasiveness. Activation of IGF-R results in its Mdm2-dependent ubiquitination and degradation followed by MAPK signaling. IGF-1R ubiquitination by Mdm2 is mediated by scaffolding protein  $\beta$ -arrestin 1. The tumor suppressor p53 pathway is activated in damaged cells causing growth arrest and if necessary, apoptosis and senescence. In normal conditions p53 is inactivated by Mdm2. Activation of extracellular pro-survival signaling has been shown to inhibit p53 activity through  $\beta$ -arrestin 1. Thus the same Mdm2/ $\beta$ -arrestin 1 system regulates two important pathways involved in cancer.

The aim of this thesis was to investigate in detail the IGF-1R/ $\beta$ -arrestin/Mdm2/p53 axis and explore the potential use of its components as anti-cancer therapeutic targets.

In **Paper I** we analysed the molecular interplay between p53 and IGF-1R through Mdm2. We tested the effect of p53/Mdm2 disruption on IGF-1R using a panel of melanoma cells and the p53-rectivator Nutlin-3. Disruption of the p53/Mdm2 interaction by Nutlin-3 increased the IGF-1R/Mdm2 interaction, followed by IGF-1R degradation and MAPK activation. This resulted in reduced cell proliferation and invasion and had a two-step effect on cell migration, demonstrating that modulation of the p53/Mdm2/IGF-1R axis is a potential anti-cancer therapeutic strategy.

In **paper II** we focused on the role of  $\beta$ -arrestin isoforms in the p53/Mdm2/IGF-1R axis. By modulating levels of  $\beta$ -arrestin 1 or 2 we identified opposing roles of isoforms on IGF-1R degradation, signaling and p53 pathway. We revealed a higher affinity of ligand-free IGF-1R for  $\beta$ -arrestin 2, and ligand occupied receptor - for  $\beta$ -arrestin 1. Antagonism between isoforms was also observed on biological effects with  $\beta$ -arrestin 2 causing cell cycle arrest and inhibiting IGF-1 response and cell viability, and  $\beta$ -arrestin 1 acting in the opposite direction. Thus we identified the  $\beta$ -arrestin 1/2 system as a second potential drug target within the p53/Mdm2/IGF-1R axis.

In **paper III** we studied the possibility of co-targeting the p53/Mdm2/IGF-1R and the MAPK pathway in melanoma cell lines. We combined MEK inhibitors with 1) balanced IGF-1R inhibition by siRNA; 2) biased IGF-1R inhibition by Nutlin-3, inducing transient MAPK; and 3) biased IGF-1R inhibition by antibody CP, inducing prolonged MAPK. We identified strong synergy between Nutlin-3 and MEK inhibitors. This combination of specific biased IGF-1R inhibition with MEK inhibitors is the first rational anti-cancer strategy identified in this thesis.

In **paper IV** we investigated the possibility of co-targeting the  $\beta$ -arrestin system with the DNA-damage inducing drug dacarbazine. By modulating the level of  $\beta$ -arrestin isoforms in melanoma cell lines we demonstrated that both  $\beta$ -arrestin 1 inhibition and  $\beta$ -arrestin 2 overexpression synergize with dacarbazine. This study revealed the second rational anti-cancer strategy of this project.

To sum up, our findings demonstrate that the p53/Mdm2/IGF-1R axis is a potential target for anti-cancer therapy. However, optimal effects can be achieved only through accurate modulation of multiple pathways regulated by the axis.

## LIST OF SCIENTIFIC PAPERS

- I. Worrall C, **Suleymanova N**, Crudden C, Trocoli Drakensjö I, Candrea E, Nedelcu D, Takahashi S-I, Girnita L and Girnita A. Unbalancing p53/Mdm2/IGF-1R axis by Mdm2 activation restrains the IGF-1-dependent invasive phenotype of skin melanoma *Oncogene* 2017 Jun 8;36(23):3274-3286; doi: 10.1038/onc.2016.472  
PMID: 28092675
- II. **Suleymanova N**, Crudden C, Shibano T, Worrall C, Oprea I, Tica A, Calin GA, Girnita A and Girnita L. Functional antagonism of  $\beta$ -arrestin isoforms balance IGF-1R expression and signaling with distinct cancer-related biological outcomes *Oncogene* doi: 10.1038/onc.2017.179; Epub ahead of print 5 June 2017  
PMID: 28581517
- III. **Suleymanova N\***, Crudden C\*, Worrall C, Dricu A, Girnita A and Girnita L. Enhanced response of melanoma cells to MEK inhibitors following unbiased IGF-1R down-regulation  
*Oncotarget* 2017 In press
- IV. **Suleymanova N**, Crudden C, Gebhard B, Girnita L, Girnita A. Competing engagement of  $\beta$ -arrestin isoforms balance IGF-1R signaling and control response of melanoma cells to chemotherapy  
*Manuscript*

\* Equal contribution

## RELATED PUBLICATIONS

- I. Crudden C, Ilic M, **Suleymanova N**, Worrall C, Girnita A, Girnita L. The dichotomy of the Insulin-like growth factor 1 receptor: RTK and GPCR: friend or foe for cancer treatment?  
*Growth Horm IGF Res.* 2015 Feb;25(1):2-12.  
doi: 10.1016/j.ghir.2014.10.002. Epub 2014 Oct 28. Review.  
PMID: 25466906
- II. Worrall C, Nedelcu D, Serly J, **Suleymanova N**, Oprea I, Girnita A, Girnita L. Novel mechanisms of regulation of IGF-1R action: functional and therapeutic implications.  
*Pediatr Endocrinol Rev.* 2013 Jul;10(4):473-84. Review.  
PMID: 23957198



## CONTENTS

1. Introduction.....	1
1.1 Cancer and its hallmarks.....	1
1.2 Growth factors and tumor suppressors.....	2
1.3 Growth factor receptors and cancer.....	3
IGF-1R system.....	3
IGF-1R as an RTK/GPCR hybrid.....	6
GPCRs.....	7
GPCRs: G protein.....	7
GPCRs: GRKs.....	8
GPCRs: $\beta$ -arrestins.....	8
GPCRs: biased signalling.....	12
IGF-1R as a GPCR.....	12
IGF-1R: G protein.....	12
IGF-1R: GRKs.....	12
IGF-1R: $\beta$ -arrestins.....	12
IGF-1R: biased signalling.....	14
$\beta$ -arrestins and cancer.....	15
1.4 Tumor suppressor p53.....	16
1.5 Link between IGF-1R signaling and p53 pathway.....	17
2. Aim of the thesis.....	19
3. Results and Conclusions.....	20
Paper I.....	20
Paper II.....	22
Paper III.....	24
Paper IV.....	26
4. Discussion and future prospectives.....	28
5. Acknowledgements.....	30
6. References.....	31

## LIST OF ABBREVIATIONS

A-loop	Activation loop
ALL	Acute lymphoblastic leukemia
AP2	Adaptor protein 2
AR	Androgen receptor
ARF	ADP-ribosylation factor
AT1R	Angiotensin II type 1 receptor
ATP	Adenosine triphosphate
BAD	Bcl-2-associated death promoter
Bcl-2	B-cell lymphoma 2
Bcl-X1	B-cell lymphoma-extra large
BRCA1	Breast cancer gene 1
Cas	Crk-associated substrate
CML	Chronic myelogenous leukemia
CXCL12	C-X-C Motif Chemokine Ligand 12
CXCR	C-X-C chemokine receptor
Cytochrome C	Cytochrome complex
D1A	Dopamin 1A
DNA	Deoxyribonucleic acid
DOR	$\delta$ opioid receptor
DTIC	Dacarbazine
ECM	Extracellular matrix
EDG1	Endothelial differentiation gene 1
EDG2	Endothelial cell differentiation gene-2
EGF	Epidermal growth factor
EGFR	Epidermal growth factor receptor
ERK	Extracellular signal-regulated kinases
ETAR	Endothelin A receptor
FACS	Fluorescence-activated cell sorting
FGF	Fibroblast growth factor
FGFR	Fibroblast growth factor receptor
FPR1	Formyl peptide receptor 1
GBM	Glioblastoma
GDP	Guanosine-5'-diphosphate
GEF	Guanine exchange factor
GPCR	G-protein-coupled receptor
Grb2	Growth factor receptor-binding protein 2
GRK	G protein-coupled receptor kinase
GSK-3 $\beta$	Glycogen synthase kinase 3 $\beta$
GTP	Guanosine-5'-triphosphate
HEK	Human embryonic kidney cells
HIF-1 $\alpha$	Hypoxia-inducible factor 1 $\alpha$
IGF-1/2	Insulin-like growth factor 1/2
IGF-1R/IGF-2R	Insulin-like growth factor 1/2 receptor
IGFBP	Insulin-like growth factor-binding protein
IP	Immunoprecipitation
IR	Insulin receptor
IRS	Insulin receptor substrate

JNK3	c-Jun N-terminal kinase 3
KISS1R	KiSS1-derived peptide receptor
KO	Knockout
LPAR	Lysophosphatidic acid receptor
MAPK	Mitogen-activated protein kinase
MBM	Medulloblastoma
Mdm2	Mouse double minute 2 homolog
MEF	Mouse embryonic fibroblast
MEK	MAPK/ERK kinase
MMP	Matrix metalloproteinase
MOR	$\mu$ opioid receptor
N1	Neurotensin 1
nAChR	Nicotinic acetylcholine receptor
Nedd4	Neuronal precursor cell-expressed developmentally downregulated 4
NF- $\kappa$ B	Nuclear factor kappa-light-chain-enhancer of activated B cells
NK1R	Neurokinin-1 receptor
p125Fak	p125 focal adhesion kinase
p130Cas	p130 Crk-associated substrate
pAkt	Phospho-Akt
PAR1/2	Protease-activated receptors 1/2
PDE	Phosphodiesterase
PDGF $\beta$	Platelet-derived growth factor $\beta$
PDGFR	Platelet-derived growth factor receptors
PK1	3-phosphoinositide-dependent protein kinase 1
pERK	Phospho-ERK
PGE2	Prostaglandin E2
PI3K	Phosphoinositide 3-kinase
PIP2	Phosphatidyl-inositol bisphosphate
PIP3	Phosphatidyl-inositol trisphosphate
PKC $\epsilon$	Protein kinase C $\epsilon$
PLC $\gamma$	Phosphoinositide phospholipase C $\gamma$
PPP	Picropodophyllin
pRb	Retinoblastoma protein
PTB	Phosphotyrosine-binding domains
pVHL	von Hippel-Lindau tumor suppressor
RTK	Receptor tyrosine kinase
S412	Serine-412
SH2	Src homology region 2 domain
Shc	Src homology 2-containing protein
Sos	Son of sevenless
SREBP	Sterol regulatory element-binding protein
TK	Tyrosine Kinase
TP53	Tumor protein 53
TP $\beta$ R	Thromboxane A2 $\beta$ -type receptor
TrkA	Tropomyosin receptor kinase A
TrkB	Tropomyosin receptor kinase B
TSP1	Thrombospondin 1
T $\beta$ RIII	TGF- $\beta$ co-receptor type III
uPA	Urokinase plasminogen activator

uPAR	Urokinase plasminogen activator receptor
V2	Vasopressin 2
V2R	Vasopressin 2 receptor
VEGF	Vascular Endothelial Growth Factor
VEGFR	Vascular Endothelial Growth Factor Receptor
WB	Western blot
WT1	Wilms' tumour suppressor 1
$\alpha$ 1AR	$\alpha$ 1 adrenergic receptor
$\alpha$ 2AR	$\alpha$ 2 adrenergic receptor
$\beta$ -arr	$\beta$ -arrestin
$\beta$ 1AR	$\beta$ 1 adrenergic receptor
$\beta$ 2AR	$\beta$ 2 adrenergic receptor

# 1. INTRODUCTION

## 1.1. Cancer and its hallmarks

Cancer is one of the leading causes of morbidity worldwide, accounting for approximately 13% of all deaths each year [1]. The number of diagnosed cases of cancer increases every year, particularly in countries with an ageing population [1]. More than 30% of cancer deaths could be avoided by better control of risk factors, early diagnosis and treatment [2-5].

Cancer development is a complex and multi-step process during which initially normal cells acquire various capabilities to grow, survive and disseminate [6]. These capabilities include six core and two emerging hallmarks of cancer [6].

Core hallmarks of cancer include:

- (i) **Sustaining proliferative signaling:** In normal cells entry into a new cycle is under the control of growth factors, such as epidermal growth factor (EGF), vascular endothelial growth factor (VEGF), insulin-like growth factor 1 (IGF-1) etc [7]. In cancer cells these pathways are dysregulated by the increased production of growth factors or receptors as well as by structural changes of the receptor and/or components of its downstream signaling [8-10].
- (ii) **Evading growth suppressors:** Cancer cells also have to bypass multiple tumor suppressor pathways that limit proliferation of normal cells [11]. Tumor suppressor proteins such as retinoblastoma protein (pRb) and p53 are able to induce cell cycle arrest or apoptosis in response to various extra- and intracellular signals [12, 13]. In cancer cells these proteins can be lost or mutated and gain new carcinogenic functions [14, 15].
- (iii) **Resisting cell death:** Normal cells in stressed conditions undergo cell death via apoptosis, autophagy and necrosis through the p53 pathway [16, 17]. Tumor cells can avoid apoptosis through increased proliferative signaling or modulation of anti-apoptotic and pro-apoptotic factors [18]. However, tumor cells can also induce cell death in order to recruit carcinogenic inflammatory agents [19, 20].
- (iv) **Enabling replicative immortality:** Chromosomal DNA of cells is protected by telomeres that shorten in time increasing possibilities of chromosomal fusions, leading to cell senescence and apoptosis [21]. Cancer cells are able to conserve telomere length, by increased production of the protective enzyme telomerase [22-25].
- (v) **Inducing angiogenesis:** In normal adult tissues, angiogenesis is activated transiently in accordance with physiological needs [26]. In cancer cells it is constantly activated by dysregulation of pro-angiogenic growth factors, such as VEGF, fibroblast growth factor (FGF) and thrombospondin 1 (TSP1), sustaining a tumor's nutrient supply and providing a transport system for metastasizing cells [27-29]. Moreover, the angiogenic switch is also facilitated by infiltration of tumor by cells of the immune system [30, 31].
- (vi) **Activating invasion and metastasis:** Normally cells are attached to each other as well as to the extracellular matrix [32]. Cancer cells can lose these abilities by

expression of embryonic factors enabling epithelial-mesenchymal transition and dysregulation of adhesion factors enabling motility and invasiveness [33, 34]. Immune cells surrounding the tumor can contribute by activating proteolytic enzymes or growth factors [35, 36]

Emerging hallmarks of cancer include:

- (vii) **Reprogramming energy metabolism:** Normal cells undergo two stages of glycolysis and switch to the single stage glycolysis in anaerobic conditions [37]. Cancer cells can switch to the single stage glycolysis even in the presence of the oxygen thus possibly redirecting glycolytic intermediates towards neighboring and new cells [38-40]. This switch can be induced by activation of oncogenes or alteration of tumor suppressors [41, 42].
- (viii) **Evading immune destruction:** Cells are constantly scanned by immune cells and eliminated in case of aberrancy [43, 44]. Cancer cells acquire the ability to hide and evade immune system by producing immunosuppressive factors or recruiting immunosuppressive inflammatory cells [45-48].

Two enabling characteristics of cancer are required for the acquisition of hallmark of cancer by normal cells [6, 49]. These characteristics include:

- (i) **Genome instability and mutation:** Various alterations in the cellular genome enable acquisition of the hallmarks of cancer [50]. Mutations of certain genes involved in guarding DNA, such as p53, facilitate further acquisition of carcinogenic mutations and survival of transformed cells [25, 51].
- (ii) **Tumor promoting inflammation:** Inflammation and recruitment of the immune cells also helps to acquire hallmarks of cancer by supplying cells with various survival, growth and proangiogenic factors as well as mutagenic oxygen species [20, 52, 53].

In addition, the **tumor microenvironment** can exploit neighboring normal cells, creating beneficial surroundings for tumor growth thus acting as a contributing factor to cancer [35, 54, 55].

Various reconsideration of classification were proposed such as including such characteristics as genome instability and stress phenotypes as a separate emerging hallmarks of cancer [56].

## 1.2. Growth factors and tumor suppressors

The balance between two important biological pathways are involved in most if not all hallmarks of cancer: growth factor receptor signaling and tumor suppressor pathways [57-61].

**Growth factors** bind to their receptors, activating signaling pathways that regulate proliferation, survival and differentiation of normal cells [62-65]. These pathways become potentially oncogenic when any of their components are hyper-activated due to various genetic or epigenetic alterations [8, 66, 67].

The anti-oncogenic **tumor suppressors** negatively regulate cell proliferation and normally induce death of incompatible cells [61]. This pathway becomes oncogenic when its components

lose their activity as a result of various mutations [61, 68]. Typically both copies of tumor suppressor genes must be mutated for malignant transformation of cell [69].

### **1.3. Growth factor receptors and cancer**

Most of the growth factor receptors belong to the family of receptor tyrosine kinases (RTKs) [10, 70]. The RTK family includes 58 receptors with related structure [71, 72]. Most RTKs are monomers with extracellular, transmembranous and intracellular domains with protein kinase activity [10, 73]. Ligand binding to the extracellular domain induces RTK dimerization [10, 71, 73]. Dimerized monomers cross-phosphorylate tyrosine residues in the intracellular domains of each other, increasing the receptor-kinase activity and creating docking sites for the proteins involved in signal transduction [73, 74]. Various downstream molecules containing SH2 or PTB domains are directly or indirectly recruited to these phosphotyrosines, transmitting information to the cell's nucleus and other regulatory organelles [71, 75, 76]. The most well studied examples of activation cascades downstream of RTKs are the mitogen-activated protein kinase (MAPK) and phosphatidylinositol 3-kinases (PI3Ks) pathways [77].

Increased production or hyperactivation of different component of RTK signaling contributes to cancer development [9, 10, 36, 78]. Some of the RTKs involved in cancer development include epidermal growth factor receptors (EGFR), vascular endothelial growth factor receptors (VEGFR), platelet-derived growth factor receptors (PDGFR), fibroblast growth factor receptor (FGFR) and insulin-like growth factor receptor IGF-1R [9, 10].

The EGFRs regulate proliferation, differentiation and migration of normal cells [79]. Increased production of ligands and/or receptors as well as mutations of the tyrosine kinase domain are found in diverse types of cancers and associated with poor prognosis [10, 80].

VEGFR is mainly expressed on endothelial cells and plays a key role in vasculo-, angio- and lymphoangiogenesis [81, 82]. VEGFRs contribute to carcinogenesis by formation of blood and lymph vessels in the tumor and its microenvironment [83]. Up-regulation of VEGFR is found in various types of malignancies [84].

PDGFR normally controls proliferation and migration of mesenchymal cells [85]. Autocrine signaling maintained by this receptor has been demonstrated in many tumors [86, 87].

FGFRs are involved in organ development during embryogenesis as well as in tissue homeostasis, proliferation, inflammation and angiogenesis in adults [28, 65]. Gene amplification, chromosomal translocations, gene fusions and missense mutations of FGFRs are found in various types of cancers [28].

### **IGF-1R system**

The IGF-1R system consists of two ligands (IGF-1 and IGF-2), three cell membrane receptors (IGF-1R, IGF-2R and IR) and seven binding proteins (IGFBP-1-7) [88]. IGF-1 and IGF-2 ligands share 62% homology and are mainly secreted by hepatocytes after growth hormone stimulation [89]. IGF-1 ligand in turn inhibits the secretion of growth hormone by the pituitary glands [90]. Around 90% of circulating IGF is inactive through binding to IGFBPs [91, 92].

### *IGF-1R structure*

IGF-1R belongs to the family of insulin receptors with a structure more complex than most RTKs [73, 74, 93]. The IGF-1R gene encodes for a single precursor that is cleaved generating two  $\alpha$  and two  $\beta$  subunits [94]. Distinct subunits are heterodimerized after glycosylation further forming a tetramer stabilized by disulfide bonds ( $\beta$ - $\alpha$ - $\alpha$ - $\beta$ ) [94, 95].  $\alpha$ -subunits are entirely extracellular and contain an IGF-1 binding site, while  $\beta$  subunits span the membrane and consist of extracellular, transmembranous and intracellular parts [93, 95]. The intracellular part of the  $\beta$ -subunit is divided into a juxtamembrane domain, tyrosine kinase domain and C-terminal domain [96].


### *IGF-1R function*

In an inactive state the activation loop (A-loop) of the intracellular tyrosine kinase domain of IGF-1R behaves as a pseudo-substrate, blocking both substrate and ATP-binding sites of the receptor [97]. IGF-1 and IGF-2 binding to the  $\alpha$ -subunit of IGF-1R causes conformational changes in the receptor followed by autophosphorylation of tyrosines at positions 1131, 1135 and 1136 of the A-loop by their dimeric subunit partner [71, 97]. Initial phosphorylation of Y1135 induces a stable structure followed by phosphorylation Y1131 and Y1136 [97, 98]. These conformational changes of the A-loop allow ATP binding and activation of several other tyrosine residues including Y950 at juxtamembrane domain of the receptor [97, 98]. Activated Y950 serves as a docking site for insulin receptor substrate 1 (IRS1) and Src homology 2-containing protein (Shc) that link the activated receptor to downstream pathways activating tyrosine kinase signaling (TK signaling) of IGF-1R [97, 98]. (Figure 1)

IRS-1 binds to the regulatory p85 subunit of PI3Ks, which allows the activation of its catalytic p110 subunits [99]. Activated PI3K phosphorylates the phosphatidyl-inositol bisphosphate (PIP2) and generates phosphatidyl-inositol trisphosphate (PIP3), which interacts with serine threonine kinase Akt [95, 99]. This interaction causes translocation of the complex to the inner membrane and activation of Akt by 3-phosphoinositide-dependent protein kinase 1 (PDK1) [100, 101]. Activated Akt induces numerous cellular processes including cell cycle progression, cell migration and anti-apoptosis signals [99, 102]. (Figure 1)

Shc or IRS-1 also recruits the growth factor receptor-binding protein 2 (Grb2) that interacts with son of sevenless (SOS), a plasma membrane guanine exchange factor (GEF) for Ras protein [103, 104]. Activated Ras interacts with Raf, which phosphorylates and activates MAPK/ERK kinase (MEK), which in turn phosphorylates and activates extracellular signal-regulated kinases (ERKs) [88, 105]. This signaling cascade is called MAPK pathway [106]. Activated ERKs phosphorylate a number of cytosolic substrates and translocate to the nucleus, activating various transcription factors that regulate cellular proliferation, differentiation and transformation [107, 108]. (Figure 1)





**Figure 1.** Schematic illustration of signaling cascades downstream of IGF-1R: Activated IGF-1R recruits 1) Shc inducing MAPK pathway, which leads to the transcriptional activation of genes involved in cell proliferation, transformation and differentiation; and 2) IRS1-4 inducing PI3K pathway, involved in protein synthesis through mTOR and anti-apoptosis through inhibition of BAD.

### Biological effects

The IGF-1R axis has been shown to play important roles in neonatal and pubertal development by inducing cell proliferation, differentiation and inhibiting apoptosis [109]. It is also important for maintenance of normal cell growth, metabolism and adhesion [110, 111]. Disruption of IGF-1 and IGF-2 genes result in dwarfism, decreased viability and significant weight loss in mice, that survive, while disruption of the IGF-1R gene results in a more severe dwarfism and death shortly after birth [112]. Double disruption of both IGF-1 and IGF-1R genes results in a phenotype similar to that seen in the disruption of IGF-1R, while double disruption of IGF-2 with IGF-1R and IGF-1 with IGF-2R genes results in a further increase in dwarfism [112].

The first report of IGF-1R involvement in cancer was by Pollak *et al.* in 1987 [113]. Since then a growing body of evidence indicates critical roles for the IGF-1R axis in cancer [111]. Various components of IGF-1R axis are elevated during cancer development [111, 114, 115]. Increased level of IGF-1 and IGF-2 ligands and IGF-1R were found in various types of cancer [116-118]. Production of IGF-1 can increase via endocrine, paracrine and autocrine regulation by cancer cells [119-121]. Numerous studies also identified high levels of IGF-1R expression in tumors due to mutations causing IGF-1R gene amplification or other unknown mechanisms [122]. In addition, other components of the IGF-1R system, such as IRSs and IGFBPs can be deregulated in different tumors [123-127]. However, several studies indicate inverse correlation with level of IGF-1R and cancer progression [128-131]. For instance, it was reported that the level of IGF-1R is decreased particularly at late stage of prostate cancer development [128-131]. The precise reason behind this controversy remains to be elucidated.

Up-regulation of components of the IGF-1R axis induces tumor development through increased

cell proliferation, anti-apoptosis, transformation, migration and invasion [111, 114, 115]. IGF-1R induces cell proliferation through the activation of the MAPK pathway downstream receptor substrates IRS-1 and Shc [132-134]. However, the same pathways are responsible for IGF-1R induced cell differentiation [135]. It has been shown that one of the factors that control the switch between IGF-1R induced proliferation and differentiation is the balance between substrates: IRS-1 predominance induces proliferation and malignant transformation, while Shc predominance induces differentiation [135].

Cells lacking IGF-1R are unable to be transformed by several oncogenes including SV40 T antigen and/or activated Ha-Ras, c-Src, EGFR and bovine papillomavirus E5 [136]. On the other hand v-Src and GTPase deficient Gα13 are able to sustain transformation in the absence of IGF-1R [137, 138]. One of the explanations is the ability of v-Src and Gα13 to bypass IGF-1R requirement by tyrosine phosphorylating the IGF-1R substrate IRS-1, directly involved in transformation, however the exact mechanism of IGF-1R induced cell transformation remains to be investigated [139].

IGF-1R protects against apoptosis via the activation of PI3K, MAPK and 14.3.3/Raf-1/Nedd4 pathways that phosphorylates BAD and inhibits its heterodimerization with anti-apoptotic proteins Bcl-2 and Bcl-Xl, which in turn inhibits pro-apoptotic proteins Bak and Bax and cytochrome C release [105, 140-142]. This prevents caspase activation and sustains mitochondrial integrity [105, 108, 140-142].

It has been shown that proliferative and anti-apoptotic effects of IGF-1R require the tyrosine kinase domain with either juxtamembrane or C-terminal domains, while cell differentiation and transformation require all three domains of the receptor [137, 143, 144].

The IGF-1R system regulates cell motility and invasiveness through PI3K mediated up-regulation of p125 focal adhesion kinase (p125Fak), p130 Crk-associated substrate (p130Cas), paxillin and matrix metalloproteinases 2 and 9 (MMP-2 and 9) [145-153]. However, dual effects of IGF-1R on MMP production have been reported – stimulatory in the presence of an activated PI3K pathway and inhibitory in conditions with predominantly activated MAPK pathway [154]. Another way of increasing tumor metastasis is IGF-1 and IGF-2 induced angiogenesis through activation of hypoxia-inducible factor 1α (HIF-1α) and transactivation of other growth factors such as VEGF [155-157]. It was also demonstrated that IGF-1 can activate urokinase plasminogen activator (uPA)/uPA receptor system (uPAR) thus leading to tumor invasion [158]. Thus the IGF-1R system can regulate tumor invasion and metastasis at multiple levels [159, 160].

### **IGF-1R as an RTK/GPCR hybrid**

It has been increasingly recognized during the last decade that despite being a prototypical RTK, IGF-1R also shares all functional characteristics of G-protein-coupled receptors (GPCRs) [88]. IGF-1R was shown to utilize G-protein signaling, G protein receptor kinase (GRK)/β-arrestin induced desensitization, receptor ubiquitination, trafficking and signaling as well as the concept of biased signaling of GPCRs [88].

## GPCRs

GPCRs are the largest family of cell surface receptors, and share a similar seven-transmembranous structure and mechanism of activation [161]. In basal conditions, a GPCR is located adjacent to heterotrimeric G proteins that consist of  $G\alpha$ -GDP,  $G\beta$  and  $G\gamma$  subunits [162]. A ligand bound receptor undergoes conformational changes to activate its GEF function followed by association with the  $G\alpha$ -GDP subunit and exchange of its GDP to GTP [163, 164]. This leads to dissociation of  $G\alpha$ -GTP from the  $G\beta\gamma$  dimer, its interaction with second messengers and activation of the receptor signaling [163]. Intrinsic GTPase activity of  $G\alpha$ -subunit causes hydrolysis of GTP to GDP returning the subunit to its inactive state [165].  $G\beta\gamma$  triggers activation of GRKs, which phosphorylate the C-terminus of the receptor, recruiting  $\beta$ -arrestins [166].  $\beta$ -arrestin binding uncouples the receptor from the  $G\alpha$  subunit leading to degradation of second messengers and further signal cessation [167]. However, there are also alternative models suggesting that certain GPCR/G protein complexes pre-exist and persist, but rearrange after receptor activation [168-174].

### GPCRs: G protein

According to the classical model, when the ligand binds to a GPCR triggering dissociation of activated  $G\alpha$ -GTP from  $G\beta\gamma$  subunits, both partners further activate their specific effector enzymes [175, 176]. There are four different G proteins based on their distinct  $\alpha$  subunit:  $G\alpha_s\beta\gamma$ ,  $G\alpha_{i/o}\beta\gamma$ ,  $G\alpha_{q/11}\beta\gamma$  and  $G\alpha_{12/13}\beta\gamma$  [177, 178]. G protein specificity of GPCRs plays an important role in determination of the proper signaling downstream of the receptor [179-181]. Table 1 illustrates the involvement of different G-protein families in various GPCRs.

<b><math>G\alpha</math> family</b>	<b>Signal transduction</b>	<b>GPCRs</b>	<b>References</b>
$G\alpha_s$ family	Stimulation of adenylyl cyclase	$\beta$ 1AR, $\beta$ 2AR, V2R	[177, 178, 182]
$G\alpha_i$ family	Inhibition of adenylyl cyclase and regulation of ion channels and activation of phosphodiesterase 6	$\alpha$ 2AR, $\beta$ 1AR, $\beta$ 2AR, PAR1	[177, 178, 183, 184]
$G\alpha_q$ family	Activation of phospholipase C	$\alpha$ 1AR, AT1R, PAR1, PAR2	[177, 185-187]
$G\alpha_{12/13}$ family	Activation of Rho GTPases	PAR1, PAR2	[177, 178, 188]

**Table 1.** *G-protein families, their signal transduction pathways and examples of coupled GPCRs.*

Some GPCRs interact with several classes of  $G\alpha$  proteins activating different signaling pathways [189, 190].

Additionally  $G\beta\gamma$  utilizes five different  $G\beta$  subtypes and eleven  $G\gamma$  subtypes that also regulate effectors such as adenylyl cyclase, ion channels, PI3K, phospholipase C [175, 191]. Dissociated  $G\beta\gamma$  is also isoprenylated interacting with the plasma membrane and recruiting GRKs close to the activated receptor [192, 193].

## GPCRs: GRKs

GRKs are receptor associated serine/threonine kinases that phosphorylate serine or threonine residues within the C-terminus of the activated GPCRs, thus reducing G protein mediated signaling [194]. There are seven GRK isoforms divided into three groups: the visual GRK1 and GRK7 subfamily, the GRK2 and GRK3 subfamily, and the GRK4, GRK5 and GRK6 subfamily [195-197]. Visual GRKs are expressed only in the retina, GRK4 is expressed only in the testis while all the other GRKs are expressed ubiquitously [195]. GRK isoforms differently phosphorylate GPCRs leading to distinct types of receptor signaling, trafficking and biological outcomes [195-197]. The term GRK phosphorylation “barcode” is used to describe different GRK-dependent phosphorylation patterns of GPCRs [195]. The fact that GRK2 KO mice (-/-) die at the embryonic stage while GRK3 and GRK5 KO mice are viable confirms non-identical roles of GRK isoforms [198, 199]. It has been shown that GRK2 and 3 play major roles in desensitization of  $\beta$ -2 adrenergic receptor ( $\beta$ 2AR) and angiotensin II type 1 receptor (AT1R), while GRK 5 and 6 are involved in their  $\beta$ -arrestin mediated signaling [200-204]. However, it was also demonstrated that GRK 2 predominantly mediates  $\beta$ 2AR desensitization in cardiac tissue, while in uterine smooth muscle this function carried out by GRK 6 [205]. Thus GPCR/GRK interaction differs depending on class of the receptor, biological function and organ specificity [195-197].

## GPCRs: $\beta$ -arrestins

Arrestins were firstly discovered as proteins that together with GRKs attenuate receptor signaling, a process called receptor *desensitization* [199]. There are four different arrestin isoforms: arrestins 1 and 4 regulate photoreceptors and are expressed predominantly in the retina, and arrestin 2 and 3 (called  $\beta$ -arrestin 1 and 2) regulate most GPCRs and are expressed ubiquitously [206]. Both  $\beta$ -arrestins are elongated molecules with two domains: N-domain and C-domain with a connecting C-terminal tail [206]. A polar core between the two domains is mainly responsible for  $\beta$ -arrestin binding to the phosphorylated receptor, though both N- and C-domain contain receptor-binding elements [207-209].  $\beta$ -arrestin isoforms have identical N-terminus and different C-terminus, which promotes differences in receptor binding [210]. Both  $\beta$ -arrestin 1 KO ( $\beta$ arr1<sup>-/-</sup>) and  $\beta$ -arrestin 2 KO ( $\beta$ arr2<sup>-/-</sup>) mice survive while double KO mice ( $\beta$ arr1<sup>-/-</sup> and  $\beta$ arr2<sup>-/-</sup>) show neonatal lethality, confirming that at least one isoform is important for the vital function of  $\beta$ -arrestins during development [211]. Thus  $\beta$ -arrestins have distinct as well as overlapping biological functions and are able to replace each other in specific conditions [211].

Although firstly discovered as proteins mediating receptor desensitization,  $\beta$ -arrestins are now recognized as mediators of two other important functions of the receptor: i) in targeting a receptor to endocytic machinery followed by its internalization and degradation or recycling called *receptor trafficking*; and ii) activation of alternative G-protein independent receptor signaling called  *$\beta$ -arrestin (or  $\beta$ -arrestin-dependent) signaling* [210].

### *GPCR desensitization*

$\beta$ -arrestins are presumably “phosphosensors” that bind to activated and phosphorylated receptors and induce post-translational changes to the receptor [212]. The altered receptor uncouples an activated  $G\alpha$  subunit, which leads to signal cessation and receptor endocytosis [212]. In addition to that,  $\beta$ -arrestins can recruit phosphodiesterase (PDE) enzymes and induce degradation of  $G\alpha$ -GTP activated second messengers [167]. However, it was demonstrated recently that certain receptors can couple to G proteins even after  $\beta$ -arrestin binding resulting in formation of the receptor/ $\beta$ -arrestin/G protein complex, giving rise to sustained G protein signaling [172, 173].

$\beta$ -arrestin 1 KO mice (-/-) survive but exhibit decreased  $\beta$ 2AR signal cessation, suggesting that  $\beta$ -arrestin 1 plays an important role in  $\beta$ 2AR desensitization [213]. However, it was shown that *in vitro* silencing of  $\beta$ -arrestin 2 affects  $\beta$ 2AR desensitization even more than silencing of  $\beta$ -arrestin 1 [214]. Testing  $\beta$ 2AR signaling in mouse embryo fibroblasts (MEFs) lacking either or both  $\beta$ -arrestins indicated that the role of isoforms in  $\beta$ 2AR desensitization is interchangeable [199]. It has been identified that both  $\beta$ -arrestin isoforms are equally involved in desensitization of AT1R [215]. Another example is protease-activated receptors 1 and 2 (PAR1, PAR2) desensitized via both  $\beta$ -arrestins, with  $\beta$ -arrestin 1 initiating the process and thus playing predominant role [216-223].

### *GPCR trafficking*

Receptor trafficking is another key process that controls the duration of its signaling [224-226].  $\beta$ -arrestins bound to the phosphorylated receptor are able to interact with elements of the endocytic machinery such as clathrin and AP2 resulting in receptor internalization [224, 227, 228]. An internalized receptor is dephosphorylated and either recycled back to the cell surface or degraded by proteasome and lysosomes depending on the strength of the receptor/ $\beta$ -arrestin binding [224-229]. Two classes of GPCRs have been identified based on the strength of this interaction [219, 227]. Class A receptors preferentially and transiently bind to  $\beta$ -arrestin 2 followed by fast dissociation of the complex and internalization of the receptor through clathrin-coated pits [219, 227]. This class of receptor includes  $\beta$ 2AR,  $\alpha$ 1 adrenergic ( $\alpha$ 1AR),  $\mu$  opioid (MOR) and dopamine 1A (D1AR) receptors [219, 227]. Class B receptors stably bind to both  $\beta$ -arrestin isoforms followed by internalization of the receptor together with  $\beta$ -arrestins [219, 227]. This group of receptors includes AT1R, neurotensin 1 (N1) and vasopressin (V2) receptors [219, 227]. Class A receptors preferentially induce receptor recycling while class B receptors induce receptor degradation [219, 227]. The switch of C-terminal tail of the receptor changes their class affinity, indicating that the C-terminus determines class specificity [219].

### *GPCRs and $\beta$ -arrestin signaling*

The third consequence of  $\beta$ -arrestin binding to GPCRs is activation of G-protein independent  $\beta$ -arrestin signaling through interaction of  $\beta$ -arrestins with a variety of cellular molecules [210]. Receptor bound  $\beta$ -arrestins can scaffold multiple signaling molecules thus transducing signals to the MAPK, PI3K, NF- $\kappa$ B, p53 and other pathways [210, 224, 230]. This results in a variety of biological outcomes such as cell proliferation, anti-apoptosis, migration and invasion [230]. Interestingly the same pathway activated through G-protein or  $\beta$ -arrestin signaling can lead to

distinct biological outcomes [231]. For instance,  $\beta$ -arrestin activated ERK1/2 is mostly localized in the cytoplasm and interacts with cytosolic substrates, whereas G-protein activated ERK1/2 moves to the nucleus and acts as a transcription factor [232, 233]. Thus it is not surprising that  $\beta$ -arrestin activated ERK1/2 regulates proteins involved in cell motility and chemotaxis, rather than transcription activity and subsequent cell proliferation [234].

Receptor class specificity also influences the length of  $\beta$ -arrestin induced signaling: transient interaction of Class A receptors with  $\beta$ -arrestin 2 activates short  $\beta$ -arrestin signaling, while stable interaction of Class B receptors with both  $\beta$ -arrestins leads to a stable  $\beta$ -arrestin signaling [224]. However, there are exceptions from the classical pattern of  $\beta$ -arrestin mediated receptor trafficking and signaling. It has been demonstrated that PAR2 receptor is degraded faster through interaction with  $\beta$ -arrestin 1, while  $\beta$ -arrestin 2/PAR2 interaction leads to more stable association and signaling [223].

In addition regardless of the class specificity  $\beta$ -arrestin isoforms can play co-dependent or reciprocal role in regulation of the  $\beta$ -arrestin induced signaling [218, 224]. Furthermore reciprocal regulation can appear as a positive input of  $\beta$ -arrestin 1 and negative by  $\beta$ -arrestin 2 or vice versa [218].

Distinct roles of  $\beta$ -arrestin isoforms in receptor desensitization, endocytosis and signaling are partially explained by recently identified specific conformations of  $\beta$ -arrestin/GPCR interaction. It has been shown that interaction of  $\beta$ -arrestins with the C-terminus of GPCRs mediate receptor internalization and signaling, while additional interaction of  $\beta$ -arrestins with the receptor transmembrane core mediates desensitization [172, 173]. However,  $\beta$ -arrestins/GPCR interactions that discriminate the roles of isoforms in receptor endocytosis from signaling remain to be investigated.

#### *Other functions of $\beta$ -arrestins*

It is important to mention that  $\beta$ -arrestin isoforms also specifically regulate functions other than desensitization, trafficking and  $\beta$ -arrestin signaling of the receptor [235, 236].  $\beta$ -arrestin 1 can move to the nucleus in response to receptor activation, enhancing histone acetylation and gene transcription, while  $\beta$ -arrestin 2 is involved in nuclear export of Mdm2 and JNK3 kinases [237-240]. It has been suggested that these functions are controlled by non-identical C-terminal regions of the isoforms [237-240].

#### *Posttranslational modifications of $\beta$ -arrestins*

Various posttranslational modifications of  $\beta$ -arrestins such as ubiquitination and phosphorylation regulate GPCR functions [227, 241]. Ubiquitination pattern of  $\beta$ -arrestins correlates with GPCR class specificity: transient  $\beta$ -arrestin ubiquitination with class A and persistent  $\beta$ -arrestin ubiquitination with class B [227, 242]. The state of  $\beta$ -arrestin phosphorylation defines receptor endocytosis for example, dephosphorylation of serine residue 412 of  $\beta$ -arrestin 1 increases its affinity for clathrins [243-245]. Oligomerization state also controls  $\beta$ -arrestins activity [246].  $\beta$ -arrestin 1 or 2 monomers are ubiquitinated by Mdm2 and involved in receptor desensitization as well as transcriptional activity of  $\beta$ -arrestin 1 [246, 247]. However,  $\beta$ -arrestin 2 oligomers are involved in the nuclear export of Mdm2 [246-248].


### Different models of GPCR regulation by $\beta$ -arrestin 1 and 2 isoforms

The  $\beta$ 2AR belongs to the class A receptors and its C-terminus phosphorylation leads to the transient binding of  $\beta$ -arrestin 2 followed by receptor internalization and recycling [219]. However, both  $\beta$ -arrestin isoforms can induce arrestin dependent ERK activation of  $\beta$ 2AR [204]. On the other hand, AT1R belongs to class B receptors and its C-tail phosphorylation recruits both  $\beta$ -arrestins that stably bind to the receptor [219]. However, the  $\beta$ -arrestin isoforms play a reciprocal role in activation of signaling:  $\beta$ -arrestin 1 induces, while  $\beta$ -arrestin 2 inhibits the arrestin signaling of AT1R [217]. Distinct roles of  $\beta$ -arrestin isoforms in endocytosis and signaling of different GPCRs are shown in Table 2.

GPCR	Desensitization	Trafficking	Signaling	References
$\beta$ 2AR	$\beta$ -arr2> $\beta$ -arr1	$\beta$ -arr2> $\beta$ -arr1	$\beta$ -arr1 $\uparrow$ / $\beta$ -arr2 $\uparrow$	[204, 218, 219]
AT1R	$\beta$ -arr1/ $\beta$ -arr2	$\beta$ -arr1/ $\beta$ -arr2	$\beta$ -arr1 $\downarrow$ / $\beta$ -arr2 $\uparrow$	[218, 219, 232]
PAR1	$\beta$ -arr1> $\beta$ -arr2	Not essential	$\beta$ -arr1 $\uparrow$ / $\beta$ -arr2 $\downarrow$	[216-220]
PAR2	$\beta$ -arr1> $\beta$ -arr2	$\beta$ -arr1/ $\beta$ -arr2	$\beta$ -arr1 $\uparrow$ / $\beta$ -arr2 $\uparrow$	[218, 219, 221-223]

**Table 2.** Role of  $\beta$ -arrestin isoforms in GPCRs desensitization, endocytosis and signaling.  
 $\uparrow$ - stimulatory role;  $\downarrow$ - inhibitory role; > - more potent regulation.

Simplified models of G-protein, GRK and  $\beta$ -arrestin isoform interactions with the most extensively studied  $\beta$ 2AR and AT1R GPCRs are illustrated in Figure 2.



**Figure 2.** Schematic illustration of the roles of G proteins, GRKs and  $\beta$ -arrestins in  $\beta$ 2AR and AT1R signaling. a)  $\beta$ 2AR activation is followed by interaction with  $G\alpha_s\beta\gamma$ , phosphorylation by GRK2/6 and  $\beta$ -arrestin1/2 binding; b) AT1R activation followed by interaction with  $G\alpha_q\beta\gamma$ , phosphorylation by GRK2/6 and  $\beta$ -arrestin1/2 binding. Distinct effects of  $\beta$ -arrestins on desensitization, downregulation and signaling are shown with red lines. Arrowed lines indicate stimulatory and blunted lines - inhibitory effects. Transparent lines indicate weaker effects.

## **GPCRs: biased signaling**

Since the discovery of multiple pathways activating downstream of the stimulated receptor, the term biased agonism was introduced [249]. The concept of biased agonism expanded particularly after the discovery of  $\beta$ -arrestin induced signaling of GPCRs [226, 250]. This concept defines the ability of the ligand to induce and/or receptor to acquire conformation, causing activation of only specific pathways downstream of that receptor [189, 190, 251, 252]. Various ligands have been described that specifically inhibit the G protein while activating the  $\beta$ -arrestin mediated signaling pathways or vice versa [253]. The possibility to activate specific pathways could be widely applied in the drug discovery field in order to minimize side effects [254].

## **IGF-1R as a GPCR**

### **IGF-1R: G protein**

It has been demonstrated that various RTKs including the IR, IGF-1R, EGFR, PDGFR and TrkA can utilize G-proteins for the activation of their signaling [255-261].

It was shown that IGF-1R also couples to  $G\alpha_i$  subunit activating MAPK pathway that can be blocked by compounds inhibiting G-proteins [262, 263]. In a basal state,  $G\alpha_i$  and  $G\beta\gamma$  both associate with IGF-1R [264]. IGF-1 stimulation releases  $G\beta\gamma$  subunit but increases the association of IGF-1R with the  $G\alpha_i$  subunit, thus also utilizing G proteins signaling [264]. Importantly, IGF-1R induced G-protein activation can result from direct IGF-1R/G-protein interaction as well as through transactivation of GPCRs [262, 264-266].

### **IGF-1R: GRKs**

Involvement of GRKs in desensitization of various RTKs was also reported [258, 259]. Activation of EGFR, PDGFR and TrkA were shown to translocate GRK2 to the plasma membrane resulting in activation of MAPK pathway [258, 267, 268].

Activated IGF-1R recruits GRKs that phosphorylate serine residues at its C-terminus [269]. There is a contrasting effect between GRK2 and GRK6 on IGF-1R function [269]. Both GRKs interact with IGF-1R, however GRK2 phosphorylates IGF-1R serine residue 1248 and induces transient  $\beta$ -arrestin 1/IGF-1R binding, while GRK6 phosphorylates serine residue 1291, inducing stable interaction of  $\beta$ -arrestin 1 with IGF-1R [269].

### **IGF-1R: $\beta$ -arrestins**

Both  $\beta$ -arrestin isoforms were shown to interact with the IGF-1R [264, 270, 271]. It has also been demonstrated that  $\beta$ -arrestin 1 interacts with the C-terminus of IGF-1R transiently binding to serine residue 1248 and stably binding to serine residue 1291 whereas the site of  $\beta$ -arrestin 2 interactions with IGF-1R has not yet been described [272, 273].

### *IGF-1R tyrosine kinase inactivation ("desensitization")*

One described mechanism of termination of IGF-1R TK signaling is the dissociation of IRS-1 from the receptor followed by Mdm2 dependent ubiquitination and degradation of IRS-1 [274].



It was shown that  $\beta$ -arrestin 1 competes with IRS-1 for Mdm2 binding and IGF-1/insulin treatment downregulate  $\beta$ -arrestin 1 thus enhancing IRS-1/2 degradation and signal cessation [274-278]. On the other hand,  $\beta$ -arrestin 1 is directly involved in IGF-1R degradation thus also contributes to the TK signaling cessation [272].

#### *IGF-1R trafficking*

The role of the  $\beta$ -arrestin system in the downregulation of IGF-1R and other RTKs such as EGFR, PDGFR and IGF-1R has been reported [257, 272, 279].  $\beta$ -arrestin 1 is recruited to the EGFR upon ligand stimulation, inducing receptor endocytosis [264, 277, 279]. Residues 319-418 of the C terminus of  $\beta$ -arrestin 1 are involved in EGFR/ $\beta$ -arrestin 1 interaction with clathrin, indicating the importance of  $\beta$ -arrestin 1 dephosphorylation [264, 277, 279].  $\beta$ -arrestin 1 also mediates interaction of PDGF $\beta$  with the GPCR EDG1 leading to internalization of PDGFR/EDG1 complex through clathrin-coated pits [257, 280]. Ligand binding to TrkA also recruits  $\beta$ -arrestin 1 to the receptor leading to the clathrin mediated receptor endocytosis [258, 281]. It has been suggested that TrkA/ $\beta$ -arrestin 1 association is also mediated through GPCR EDG2 [258, 281]. However, the role of the  $\beta$ -arrestin 2 isoform in RTK internalization and signaling is mostly unexplored.

It is demonstrated that clathrin dependent internalization of IGF-1R is mediated by both  $\beta$ -arrestin isoforms [270]. Dephosphorylation of C-terminal serine 412 residue of  $\beta$ -arrestin 1 is a prerequisite for this internalization [270]. The ubiquitin ligase Mdm2 is recruited to activated IGF-1R through  $\beta$ -arrestin 1 leading to receptor degradation [273, 282]. Both  $\beta$ -arrestin isoforms were shown to induce IGF-1R ubiquitination and downregulation with  $\beta$ -arrestin 1 being more potent than  $\beta$ -arrestin 2 [272]. The mechanism and biological outcomes of this difference have not yet been investigated.

#### *IGF-1R and $\beta$ -arrestin induced signaling (TK independent signaling)*

Despite indication of  $\beta$ -arrestin 1 involvement in ligand-induced EGFR downregulation, no studies have shown the effect of  $\beta$ -arrestins on EGFR mediated MAPK signaling [264, 277, 279, 283]. However, it has been shown that PGE2 induced EGFR transactivation followed by Akt signaling is mediated through  $\beta$ -arrestin 1 [284]. It has been demonstrated that ligand binding to TrkA recruits  $\beta$ -arrestin 1 thus mediating activation of MAPK signaling [281]. The specific role of  $\beta$ -arrestin isoforms on PDGF and FGFR signaling remains unknown [257, 280].

Ligand dependent interaction of IR with  $\beta$ -arrestin 2 was shown to activate Akt through Src thus contributing to insulin PI3K signaling [285].  $\beta$ -arrestin 1 was not shown to affect insulin induced metabolic or mitogenic signaling pathways of IR [264, 274-277].

Recruitment of  $\beta$ -arrestin 1 to IGF-1R has been shown to mediate IGF-1 induced activation of MAPK and PI3K signaling [264, 270, 271, 286]. Activation of MAPK requires  $\beta$ -arrestin 1 dephosphorylation, its association with clathrin as well as recruitment of Mdm2 [270, 282]. The mechanism of IGF-1R/ $\beta$ -arrestin 1 mediated PI3K activation is not fully investigated [286].

## IGF-1R: biased signaling

Since the discovery of the GPCR properties of RTKs, the paradigm of biased agonism was also introduced to the field of RTKs [287-289].


Different extracellular domains of RTKs responsible for activation of distinct signaling pathways have been identified [290-296]. Various alterations of ligands, presence of co-receptors and certain intracellular proteins were shown to induce allosteric modulations of RTKs [297-301]. For instance, PKC $\epsilon$  was shown to interact with the cytosolic domain of EGFR inhibiting PLC $\gamma$  but not ERK and Akt signaling and altering sensitivity to EGFR inhibitors [297, 298]. All these studies offer new possibilities to target RTKs with various biased agonists for better therapeutic outcomes [292-296].

A limited number of biased agonists have been described for RTKs [287]. It has been demonstrated that a specific inhibitor of FGFR, SSR128129E, binds to the extracellular domain of the receptor changing its conformation and inhibiting MAPK but not phospholipase C activation [302, 303]. A compound interacting with the juxtamembrane domain of TrkA was shown to inhibit receptor inducing its dimerization and biased signaling [304].

Biased signaling of IGF-1R was first described when the kinase inhibitor cyclolignan picropodophyllin (PPP) was shown to induce IGF-1R downregulation and biased ERK signaling [305]. Similarly, the antimicrobial peptide LL-37 has been demonstrated to induce IGF-1R degradation with activation of biased signaling [306]. The IGF-1R antibody figitumumab (CP-751871 or CP), previously described as a receptor antagonist also maintains agonist like properties, it induces IGF-1R downregulation and activates  $\beta$ -arrestin 1-dependent ERK signaling [307]. Since the discovery of these properties, it becomes important to test the effect of all IGF-1R targeting molecules on all pathways downstream of IGF-1R.

To sum up, all main characteristics of GPCR activation and desensitization were detected at the IGF-1R: 1) IGF-1 binding activates the G protein signaling cascade, as well as the classical tyrosine kinase signaling [264]; 2) IGF-1R phosphorylation by GRK2 or GRK6 at C-terminal serine residues allows  $\beta$ -arrestin binding to the receptor [269]; 3) the GRK phosphorylated receptor undergoes ubiquitination and degradation or recycling to the cell surface [269, 272]; 4)  $\beta$ -arrestin recruitment leads to cessation of IGF-1R signaling [272]; 5)  $\beta$ -arrestin binding activates a second wave of IGF-1R signaling called  $\beta$ -arrestin dependent signaling [282]; 6) IGF-1R targeting molecules can be classified based on their biased agonistic properties [88]. Thus it has been proposed to classify the IGF-1R as a functional hybrid of both RTK and GPCR [88].

Simplified overview of the previously known roles for  $\beta$ -arrestins in IGF-1R activation and signaling is provided in Figure 3.



**Figure 3.** a) Absence of the ligand determines absence of the TK activity as well as IGF-1R/ $\beta$ -arrestin interaction and  $\beta$ -arrestin signaling; b) IGF-1 binding leads to TK activation, followed by GRK phosphorylation and recruitment of  $\beta$ -arrestin 1 and Mdm2 leading to receptor degradation and ERK signaling; c) CP binding inhibits the TK activity of the receptor but recruits  $\beta$ -arrestin 1 leading to receptor degradation and ERK signaling.

### $\beta$ -arrestins and cancer

$\beta$ -arrestin signaling of GPCRs and RTKs regulates multiple cancer inducing cellular functions such as cell proliferation, migration, invasion, anti-apoptotic signaling and cell cycle progression [218, 222, 224, 226, 234, 236, 284, 308-314].  $\beta$ -arrestins activate these functions by scaffolding various cytosolic proteins as well as directly regulating expression of genes [218, 224, 226, 234, 237, 315, 316]. Both  $\beta$ -arrestins are involved in different cancer related receptor functions [236, 314].

Overexpression of  $\beta$ -arrestin 1 in transgenic mice increases VEGF induced MMP-9 activity and cell growth [317]. Stimulation of various GPCRs (FPR1, AT1R, V2R, and CXCR2) induces apoptosis in  $\beta$ -arrestins 1 and 2 knockout mouse embryonic fibroblasts (MEFs) while expression of  $\beta$ -arrestins rescues this effect [309].

$\beta$ -arrestins mediate neurokinin-1 receptor (NK1R) induced ERK activation through Src, leading to cell proliferation and anti-apoptosis [318]. IGF-1 activation causes a  $\beta$ -arrestin 1 mediated increase of pERK1/2 and pAkt leading to cell proliferation and anti-apoptosis [282, 286]. The C-terminal domain of  $\beta$ -arrestin 2 plays specific role in anti-apoptotic signaling through adaptor protein AP2 [319, 320].  $\beta$ -arrestin 2 mediates AT1R induced anti-apoptotic signaling through MAPK and PI3K pathway by phosphorylation of apoptotic protein BAD [321].  $\beta$ -arrestin 2 also attenuates GSK-3 $\beta$  apoptotic signaling of Toll-like receptor 4 [322]. Similarly,  $\beta$ -arrestin 2 inhibits resveratrol induced GSK-3 $\beta$  mediated apoptosis of endometrial cancer cells [323]. CXCR4 induced cell migration is mediated by  $\beta$ -arrestin 2 activated MAPK pathways [324, 325]. However,  $\beta$ 2AR induced dephosphorylation of  $\beta$ -arrestin 2 promotes apoptosis after UV exposure via stabilization of I $\kappa$ B $\alpha$ /NF- $\kappa$ B association [326]. In addition, activated  $\delta$  opioid receptor (DOR) recruits Mdm2 through  $\beta$ -arrestin 2 thus inhibiting Mdm2/p53 interaction and increasing p53-mediated apoptosis [327].

Specific involvement of  $\beta$ -arrestin isoforms in various types of cancer summarized in Table 3.

Cancer	Isoform	Mechanism of involvement	Ref.
Colon cancer	$\beta$ -arrestin 1	Mediates PGE2 induced EGFR transactivation	[284]
	$\beta$ -arrestin 2	Mediates Wnt signaling	[313]
Bladder cancer	$\beta$ -arrestin 1	Not involved in TP $\beta$ R induced malignancy	[328]
	$\beta$ -arrestin 2	Mediates TP $\beta$ R signaling	[329]
Blood cancers	$\beta$ -arrestin 1	Elevated in ALL, inhibits Notch1 signaling	[330]
	$\beta$ -arrestin 2	Involved in CML, mediates Wnt signaling	[310]
Prostate cancer	$\beta$ -arrestin 1	Mediates Wnt signaling	[331]
	$\beta$ -arrestin 2	Downregulates AR and TP $\beta$ RIII; Mediates $\beta$ 2AR induced c-Src activation	[332-335]
Pancreatic cancer	$\beta$ -arrestin 1	N/D	
	$\beta$ -arrestin 2	Mediates CXCR4/7 induced MAPK signaling; Possibly mediates increased Shh signaling	[336-339]
Brain tumors	$\beta$ -arrestin 1	Low phospho-S412 correlated with low survival in GBM	[340]
	$\beta$ -arrestin 2	Possibly mediates increased CXCR4 signaling in MBM	[341]
NSCLC	$\beta$ -arrestin 1	Mediates PGE2 induced EGFR transactivation and nAChRs signaling; Nuclear increase correlates with poor prognosis	[308, 342, 343]
	$\beta$ -arrestin 2	Decrease correlates with poor prognosis	[312]
Breast cancer	$\beta$ -arrestin 1	Mediate LPAR and PAR2 signaling and HIF-1 $\alpha$ dependent activation of VEGF-A; Decreased with tumor progression, correlating with poor prognosis	[222, 344-346]
	$\beta$ -arrestin 2	Mediate LPAR, PAR2 and KISS1R signaling; TP $\beta$ RIII dependent inhibition of NF- $\kappa$ B; Possibly mediates CXCR4 signaling; Increased with tumor progression, correlating with poor prognosis	[222, 344, 346-349]
Ovarian cancer	$\beta$ -arrestin 1	Mediates ETAR signaling and EGFR transactivation	[311]
	$\beta$ -arrestin 2	Mediates ETAR signaling and EGFR transactivation	[311]

**Table 3.** Mechanisms of  $\beta$ -arrestin 1 and 2 involvement in various types of cancer.

#### 1.4. Tumor suppressor p53 pathway

The p53 pathway is a crucial tumor suppressor pathway that is probably inactivated in all tumors [18, 350-352]. p53 prevents cancer by maintaining genetic stability in damaged cells and eliminating incipient tumor cells [353, 354]. Oncogenic stress induces p53 phosphorylation and acetylation, which inhibits its interaction with the Mdm2 ubiquitin ligase and thereby stabilizes p53, and activates p53 as a transcription factor [353-357]. Activated p53 transcriptionally activates target genes and induces growth arrest and if necessary, apoptosis or senescence [353-356]. In an unstressed environment p53 is degraded and kept at low levels by

Mdm2 that binds to p53, mono-ubiquitinates it and exports to the cytosol where p53 is degraded in the proteasome [353-356]. Expression of Mdm2 is also transcriptionally regulated by p53 through a negative feedback loop [353-357].

The TP53 gene is mutated in around 50% of cancers [358, 359]. Most TP53 mutations are missense mutations that disrupt p53 DNA binding [358, 359]. Mutant p53 can exert a dominant negative effect on a wild type p53 [358, 359]. Another mechanism of p53 inactivation is overexpression of its natural inhibitor, the E3 ubiquitin ligase Mdm2, in cancer cells [358-361].

## **1.5. Link between IGF-1R signaling and p53 pathway**

There is a solid amount of data indicating that p53 interacts with the IGF-1 pathway 1) at the transcriptional level; 2) through regulation of metabolic processes; 3) and through convergence of the IGF-1R and p53 signaling pathways [362, 363].

### *Link at the transcriptional level*

It has been shown that wild type p53 suppresses transcription of IGF-1R, possibly as part of its apoptotic effect, however mutant p53 activates IGF-1R expression [364-366]. Wild type p53 also inhibits transcription of IGF-2 and induces transcription of the IGFBP2 and IGFBP3 genes [367-369]. In addition, p53 homologs p63 and p73 proteins also inhibit IGF-1R transcription while their mutant forms fail to do so [370]. Other tumor suppressors such as BRCA1, pVHL and WT1 were also demonstrated to reduce expression of IGF-1R [371-374].

### *Regulation of metabolic processes*

The interplay between IGF-1R and the p53 pathway is also implicated in regulation of metabolic pathways [375]. IGF-1 directly regulates glucose uptake and inhibits hepatic production of glucose [376]. However, wild type p53 inhibits glycolysis through repression of *GLUT1* and *GLUT4* genes, while mutant p53 lacks this function [377]. Wild type p53 also promotes hepatic production of glucose thus possibly preventing glycolysis utilized by cancer cells [378]. Another example is that IGF-1 stimulates expression of the transcription factor SREBP involved in sterol synthesis, while p53 inhibits SREBP and thus lipogenesis [379, 380].

### *Convergence of signaling pathways*


Mdm2 is a well-described ubiquitin ligase for both p53 and IGF-1R [273, 381]. It was demonstrated that IGF-1 signaling affects the activity of Mdm2 towards p53 [382, 383]. IGF-1R activation induces PI3K pathway leading to Akt translocation to the nucleus followed by up-regulation of transcription factors involved in cell growth and anti-apoptosis [382, 383]. However, Akt also phosphorylates Mdm2 inhibiting its association with predominantly nuclear protein ARF [382-384]. This leads to nuclear export of the Mdm2/p53 complex followed by cytoplasmic proteasomal degradation of p53 [383-386]. p53 can inhibit the PI3K pathway through activation of the inhibitor of this pathway, PTEN [387].

It was also shown that IGF-1 induces Mdm2-dependent p53 degradation in response to DNA damage through activation of MAPK pathway [384, 388-390]. In addition p53 can inhibit MAPK pathway through caspase-mediated cleavage of the ERK2 [391].

$\beta$ -arrestin 1 was shown to mediate Mdm-2 dependent IGF-1R degradation and MAPK signaling [272, 282]. Interestingly  $\beta$ -arrestin 1 also facilitates Akt induced p53 degradation by Mdm2 downstream of  $\beta$ 2AR [392]. However, the role of  $\beta$ -arrestins in IGF-1 induced Mdm2/p53 interaction has not been elucidated.

Thus IGF-1R and p53 pathway are interconnected at multiple levels and dissection of this complex interplay has a major impact on understanding of basic physiology as well as on improvement of specific anti-cancer therapies.

A simplified scheme of IGF-1R interplay with the p53 pathway is provided in Figure 4.



**Figure 4.** Schematic illustration of interplay between IGF-1R and wild type (green) and mutant (grey) p53. Dashed line indicates transcriptional regulation, solid line indicates regulation at the protein level. Arrowed lines indicate stimulatory and blunted lines indicate inhibitory effects.

## **2. AIM OF THE THESIS**

The main aim of this thesis was to investigate in detail the IGF-1R/ $\beta$ -arrestin/Mdm2/p53 axis in cancer and to explore the potential use of its components as therapeutic targets.

Paper I: To analyze the molecular interplay between p53 and IGF-1R pathways through Mdm2.

Paper II: To investigate the role of  $\beta$ -arrestin isoforms in the IGF-1R/ $\beta$ -arrestin/Mdm2/p53 axis.

Paper III: To reveal a potential of co-targeting the IGF-1R/ $\beta$ -arrestin/Mdm2/p53 axis with the MAPK pathway.

Paper IV: To reveal the potential of co-targeting the IGF-1R/ $\beta$ -arrestin/Mdm2/p53 axis with DNA damage inducing drugs.

### 3. RESULTS AND CONCLUSIONS

#### 3.1. Paper I: Unbalancing p53/Mdm2/IGF-1R axis by Mdm2 activation restrains the IGF-1-dependent invasive phenotype of skin melanoma

Background and rationale:

p53 mutations are rare in melanoma and instead p53 is often inhibited by up-regulation of the ubiquitin ligase Mdm2 [393-396]. Thus, the disruption of the Mdm2/p53 interaction could be a rational approach for melanoma treatment. The IGF-1R is another important player in melanoma cell survival and is also regulated by ubiquitin ligase Mdm2 [273, 397, 398]. As both p53 and IGF-1R are substrates of Mdm2, we investigated the impact of Mdm2/p53 disruption on function of IGF-1R in a melanoma model.

Results and conclusions:

We used the prototypical p53-reactivator Nutlin-3 to disrupt Mdm2/p53 complex in four melanoma cell lines, two with wild type (wtp53) and two with mutant p53 (mtp53).

Since Mdm2 is a ubiquitin ligase for IGF-1R, we first investigated the effect of Nutlin-3 on the levels of Mdm2 and IGF-1R. We demonstrated that in wtp53 cells, Nutlin-3 causes a dose and time dependent accumulation of p53 and Mdm2, accompanied by a decrease in the level of IGF-1R. No significant effects on the level of p53, Mdm2 or IGF-1R were observed in mtp53 cells. These data indicate that Nutlin-3 affects IGF-1R level following increase of Mdm2.

It has been demonstrated that Mdm2 induces IGF-1R ubiquitination followed by its degradation and pERK signaling in response to IGF-1 stimulation [272, 273, 282]. Thus, we next tested whether IGF-1 stimulation was essential for the Nutlin-3 mediated IGF-1R degradation. Cells were Nutlin-3 treated, serum starved, IGF-1 stimulated and monitored for changes in the level of IGF-1R using western blot (WB). We demonstrated that IGF-1R degradation only occurred in the presence of IGF-1, despite similar levels of p53 accumulation in both serum free and IGF-1 conditions. This effect was also obvious only in wtp53 cells.


We next investigated the mechanism of Nutlin-3 dependent IGF-1R downregulation. Using qPCR we excluded transcriptional origin of Nutlin-3 mediated IGF-1R decrease. Using immunoprecipitation (IP) we investigated whether IGF-1R is physically associated with and ubiquitinated by Mdm2 after Nutlin-3 treatment. In serum starved conditions recruitment of Mdm2 to the IGF-1R was very low, increasing after Nutlin-3 treatment or IGF-1 stimulation and reaching maximum after Nutlin-3 treatment followed by IGF-1 stimulation. A similar pattern was observed for ubiquitination level: Nutlin-3 or IGF-1 treatment alone increased IGF-1R ubiquitination, while double treatment increased it further. We next tested Nutlin-3 induced IGF-1R degradation in cells expressing low level of Mdm2 (SAOS) and full length (MEF wt) and C-terminus truncated ( $\Delta$ 1245) IGF-1R. In SAOS2 cells Nutlin-3 treatment did not induce either p53 or Mdm2 accumulation, while in MEF wt and  $\Delta$ 1245 it induced both. However, Nutlin-3 induced IGF-1R degradation was only observed in MEF wt cells. These data confirmed that Mdm2-mediated ubiquitination of the IGF-1R is a key mechanism of Nutlin-3 induced IGF-1R downregulation.



Since IGF-1R ubiquitination by Mdm2 also activates pERK we next tested the effect Nutlin-3 may have on IGF-1R signaling. We stimulated untreated and Nutlin-3 treated cells with IGF-1 for up to 60 minutes and observed an early and overall increase of pERK in Nutlin-3 treated samples, compared to untreated. This increase was eliminated by IGF-1R downregulation and was not observed in SAOS-2 and  $\Delta 1245$  cells. These results demonstrated that Nutlin-3 acts as a partial agonist for IGF-1R induced pERK signaling through Mdm2.

IGF-1R is a potent inducer of cell transformation, proliferation, invasion and metastasis [117, 148, 399]. Thus, we next investigated the effect of Nutlin-3 on these biological behaviours. Using a soft agar assay, cell viability assay and FACS analysis we demonstrated that Nutlin-3 treatment inhibits colony formation, slightly increases total cell number, and decreases overall IGF-1 response. Using both monolayer wound healing and transwell chamber assays we demonstrated that Nutlin-3 increased at early stage and decreased at later time points IGF-1 induced migration, particularly in wtp53 cells. However, a matrigel invasion assay and monitoring of MMP-2 activation showed that Nutlin-3 completely inhibits IGF-1 induced invasion. To sum up, Nutlin-3 inhibits IGF-1R induced cell proliferation/survival, has a two-step effect on IGF-1 induced cell migration and completely abolishes IGF-1 induced invasion.

The main findings of the study are that 1) p53/Mdm2 disruption activates IGF-1R interaction with Mdm2; 1) Nutlin-3 comprises agonistic properties towards IGF-1R induced pERK; 2) p53/Mdm2/IGF-1R axis could be used as a potential target for anti-cancer therapy.



**Figure 5.** Schematic illustration of the p53/Mdm2/IGF-1R axis in the a) absence of Nutlin-3 and IGF-1; b) presence of Nutlin-3; c) presence of both, Nutlin-3 and IGF-1.

### **3.2. Paper II: Functional antagonism of $\beta$ -arrestin isoforms balance IGF-1R expression and signaling with distinct cancer-related biological outcomes**

Background and rationale:

The roles of the two  $\beta$ -arrestins ( $\beta$ -arrests) with similar structure are well characterized in various GPCRs [224, 226]. In regards to their function at the IGF-1R, it is mainly the function of  $\beta$ -arr1 that has been described so far. It has been demonstrated that  $\beta$ -arr1 induces ligand dependent IGF-1R degradation, activates late MAPK signaling, and is able to mediate resistance to IGF-1R targeting in cancer [272, 282, 305]. However, it was also shown that both  $\beta$ -arrests can bind and ubiquitinate the IGF-1R leading to its degradation [269, 272]. Yet little is known about the specific function of  $\beta$ -arr2 on IGF-1R expression and function. In this study, we particularly focused on the role of the  $\beta$ -arr2 isoform and identified distinct roles for the two isoforms on IGF-1R system as well as on the interconnected p53 pathway.

Results and conclusion:

It has been shown that  $\beta$ -arr isoforms can differently downregulate various GPCRs [219, 225, 227]. Thus, we first studied the effect of  $\beta$ -arr2 on IGF-1R expression.  $\beta$ -arr1 knock out ( $\beta$ 1KO),  $\beta$ -arr2 knock out ( $\beta$ 2KO) and wild type (WT) MEF cells were serum starved and stimulated with IGF-1 for 0, 12 and 24 h followed by monitoring of IGF-1R degradation rate by WB. Our results indicated that IGF-1R degradation is impaired in  $\beta$ 1KO, and enhanced in  $\beta$ 2KO cells compared to WT cells. We next transfected HEK and WT MEF cells with  $\beta$ -arr1 or  $\beta$ -arr2 encoding plasmids or siRNAs and showed that  $\beta$ -arr1 up-regulation and  $\beta$ -arr2 inhibition enhances, whilst  $\beta$ -arr1 inhibition and  $\beta$ -arr2 up-regulation impairs IGF-1R degradation. Importantly, IGF-1R level was decreased in serum starved  $\beta$ -arr2 up-regulated cells prior to stimulation. This experiment demonstrated that in contrast to  $\beta$ -arr1,  $\beta$ -arr2 inhibits the rate of ligand dependent degradation of IGF-1R.

It has been also demonstrated that  $\beta$ -arr1 sustains IGF-1 induced ERK activation [282]. Therefore, we next tested the effect of  $\beta$ -arr2 on modulation of IGF-1R signaling. We serum starved and stimulated  $\beta$ 1KO,  $\beta$ 2KO and WT MEF cells with IGF-1 for 0, 2, 5, 10, 30 and 60 minutes and indicated that ERK activation lasts longer in  $\beta$ 2KO and shorter in  $\beta$ 1KO compared to WT cells. Down- or upregulation of  $\beta$ -arr2 in HEK and MEF cells confirmed that  $\beta$ -arr2 inhibition prolongs and up-regulation shortens IGF-1 activated ERK. These data indicate opposing roles for  $\beta$ -arr1 and  $\beta$ -arr2 on IGF-1-induced late ERK activation.


$\beta$ -arr1 is known to control IGF-1R degradation and signaling through Mdm2 dependent ubiquitination of the C-terminus of IGF-1R [282]. Thus, we next tested impact of Mdm2 on  $\beta$ -arr2 activity by altering  $\beta$ -arrests in MEF cells, expressing a C-terminus truncated IGF-1R ( $\Delta$ 1245), SAOS-2 cells with low endogenous and U2OS cells with high endogenous level of Mdm2. We observed a similar effect for  $\beta$ -arr2 only in U2OS cells, while in  $\Delta$ 1245 and SAOS-2 cells IGF-1R degradation and signaling was unchanged after  $\beta$ -arr2 alteration. This data indicated a role for Mdm2 and the C terminus of IGF-1R in the modulation of IGF-1R function by  $\beta$ -arr2.

We next tested whether  $\beta$ -arr2 also binds to and induces ubiquitination of the IGF-1R. We up-regulated IGF-1R and either  $\beta$ -arr1 or  $\beta$ -arr2 in U2OS cells, followed by serum starvation, IGF-1 stimulation and immunoprecipitation of the  $\beta$ -arrs. We analyzed the immunoprecipitates and demonstrated that  $\beta$ -arr2 preferentially binds to and induces ubiquitination of a ligand-free IGF-1R, while  $\beta$ -arr1 - of a ligand occupied IGF-1R.

Next, we assessed the biological outcomes of modulation of IGF-1R expression and ERK activation by  $\beta$ -arr isoforms. We modulated  $\beta$ -arr1 or 2 in U2OS and SAOS-2 cells, followed by serum starvation, IGF-1 stimulation and analysis of proliferation/survival and cell cycle distribution using PrestoBlue and FACS. We revealed increased total cell number and cell cycle progression in response to IGF-1 after  $\beta$ -arr1 up-regulation and  $\beta$ -arr2 inhibition in both cell lines.  $\beta$ -arr2 up-regulation and  $\beta$ -arr1 inhibition in contrast, reduced cell number as well as IGF-1 response causing cell cycle arrest. Interestingly, SAOS-2 were affected more and arrested in G2 phase, while U2OS cells were arrested in both G1 and G2 phases, suggesting that p53 absence facilitates a G1/S transition even without sustained proliferative signals.

We proposed that  $\beta$ -arr2 predominance decreases cell viability through  $\beta$ -arr2 mediated downregulation of IGF-1R during serum starvation and an activated p53 pathway. We confirmed both scenarios by monitoring a gradual decrease in IGF-1R level in serum free conditions and re-activation of p53 in  $\beta$ -arr2 up-regulated and  $\beta$ -arr1 inhibited cells. Thus both receptor depletion and increased p53 contributes to IGF-1 insensitivity in  $\beta$ -arr2 predominant conditions.

The main findings of this study are 1) opposing roles of  $\beta$ -arr isoforms in modulating IGF-1R function; 2) an important role of  $\beta$ -arr1/2 balance in cell cycle progression; 3) identification of anti-cancer therapeutic potential of  $\beta$ -arr1/2 system with an impact on both pro-oncogenic IGF-1R signaling and tumor suppressor p53 pathway.



**Figure 6.** Schematic illustration of prevailing manner of  $\beta$ -arr1 or  $\beta$ -arr2 interaction with p53 and IGF-1R in the a) presence of IGF-1; b) absence of IGF-1 ligand.

### **3.3. Paper III: Enhanced response of melanoma cells to MEK inhibitors following unbiased IGF-1R down-regulation**

Background and rationale:

Due to frequent mutation of RAF or RAS genes, inhibition of the downstream MAPK is a rational approach for melanoma treatment [400]. However, current MAPK inhibitors develop rapid resistance soon after administration that is associated with a more aggressive disease recurrence [401-403]. Melanoma cells acquire alternative routes to sustain proliferative signaling, including IGF-1R exploitation. Thus, co-targeting of MAPK inhibition with IGF-1R downregulation is a reasonable strategy for better outcome of treatment [404-406]. Yet IGF-1R downregulation can be associated with inappropriate activation of  $\beta$ -arr induced biased MAPK signaling [289, 305, 307, 407, 408]. In this study, we explore co-targeting of MAPK inhibition with balanced and  $\beta$ -arr signaling biased IGF-1R downregulation to aim to maximize drug sensitivity of melanoma cells.

Results and conclusion:

Since RAS and RAF mutations occur in high percentage in melanoma we first tested RAS/RAF status and sensitivity to MEK1/2 inhibitor U0126 in a panel of melanoma cell lines. We confirmed by sequencing that DFB and Mel28 cells contains BRAF mutation, while BE contains an NRAS mutation. Cells cultured in media with and without serum followed by WB analysis indicated that the highest level of basal pERK1/2 was in Mel28 and lowest in BE cells, which further increased with serum. We demonstrated a U0126 dose-dependent decrease in cell viability with a higher response in BRAF positive cells. We also demonstrated a higher sensitivity to U0126 in the absence of serum compared to presence of serum or IGF-1. Taken together, this data revealed that U0126 treatment decreased cell number in all cell lines and this effect was limited in the presence of IGF-1.


Since IGF-1R signaling is able to decrease U0126 sensitivity we next tested various strategies to down-regulate IGF-1R. As a first strategy we used siRNA mediated balanced IGF-1R downregulation that does not modify IGF-1R signaling [409, 410]. As a second strategy we downregulated IGF-1R with targeted antibodies (CP), inducing biased sustained  $\beta$ -arr mediated MAPK signaling [307]. The third strategy was to downregulate IGF-1R by Nutlin-3, inducing biased transient  $\beta$ -arr mediated MAPK signaling [411]. We identified that siRNA and CP downregulates IGF-1R in all cell lines while Nutlin-3 has effects only in DFB and to a lesser extent in BE cells. Only Nutlin-3 treated DFB cells indicated p53 re-activation. We tested the effect of treatments on IGF-1 induced proliferation of cells. We identified a decreased IGF-1 response in all siRNA, CP and Nutlin-3 treated DFB and to a lesser extend BE cells, whereas Mel28 were not affected. These results confirmed downregulation of IGF-1R and inhibition of IGF-1 response by three different strategies.

We have previously demonstrated that CP and Nutlin-3 are able to activate sustained or transient biased ERK activation [307, 411]. Therefore, we next tested the biased agonistic features of all three strategies. We treated cells with CP, Nutlin-3 and siRNAs, serum starved them and stimulated with IGF-1 for up to 60 minutes. WB analysis revealed the following patterns of pERK activation: in siRNA treated cells pERK1/2 was consistently decreased in all

cell lines compared to untreated controls; Nutlin-3 treatment induced an early increase and late decrease in the level of pERK1/2 in DFB and BE cells, compared to untreated controls; CP treatment decreased overall but sustained late pERK1/2 in all cell lines compared to untreated controls. These results indicated that all strategies decreased IGF-1R signaling, however siRNA treatment caused a balanced decrease, while Nutlin-3 and CP treatment induced biased transient and sustained pERK1/2, respectively.

We next investigated the effects of transient versus biased IGF-1R down-regulation on melanoma response to MEK1/2 inhibitor. Cells were first treated with siRNA, Nutlin-3 or CP, then with or without U0126. Results indicated a decrease in the level of IGF-1R in all cells except for Nutlin-3 treated Mel28 and an increase of p53 level only in Nutlin-3 treated DFB cells. The level of pERK was lowered in siRNA or Nutlin-3 combined with U0126 treated cells, but remained higher in CP combined with U0126 treated cells compared to control. Analysis for the total cell number indicated a 5-20% decrease after all three types of treatment. These data were combined with identical conditions of U0126 treatment alone to calculate a predicted additive response and compared with the observed response of combined treatment. We observed higher than predicted sensitivity to U0126 in all cells pre-treated with siRNA and in DFB and BE cells pre-treated with Nutlin-3, and close to predicted sensitivity to U0126 in cells pre-treated with CP. These data indicated synergy of balanced and transiently biased but not the sustained biased approach with MEK1/2 inhibition.

The main findings of this study are: 1) IGF-1R downregulation can be balanced or accompanied with transient or sustained biased signaling; 2) both balanced and transient biased IGF-1R downregulation synergizes with inhibition of MEK1/2, while sustained biased IGF-1R downregulation does not; 3) Nutlin-3 treatment offers a possible strategy to increase specificity of combined treatment.



**Figure 7.** Schematic illustration of IGF-1R and p53 activity after treatment with a) siRNAs against IGF-1R; b) p53 re-activator Nutlin-3; c) IGF-1R antibody CP

### **3.4. Paper IV: Competing engagement of $\beta$ -arrestin isoforms balance IGF-1R signaling and control response of melanoma cells to chemotherapy**

Background and rationale:

Dacarbazine (DTIC) is a chemotherapeutic agent that induces DNA damage leading to activation of p53 pathway, followed by apoptosis and cell cycle arrest. Despite poor response and due to lack of alternatives it remains the main compound used for the treatment of metastatic melanoma. The IGF-1R is an important growth factor receptor involved in proliferation and invasion of melanoma cells [133, 397-399]. It been demonstrated recently that  $\beta$ -arrests control both mitogenic IGF-1R signaling and the tumor suppressor p53 pathway [272, 282, 392, 412]. In this study we aimed to investigate the possibility of co-targeting components of  $\beta$ -arr system in order to increase sensitivity to DTIC.

Conclusion and results:

We have demonstrated previously that  $\beta$ -arr isoforms play opposing roles in IGF-1 induced cell proliferation [412]. Thus, we first investigated the effect of  $\beta$ -arr modulation in melanoma cell lines. We first verified that all three melanoma cells express IGF-1R and the two  $\beta$ -arr isoforms. We next tested the effect of  $\beta$ -arr isoforms on melanoma cell proliferation and survival by down- and upregulating either  $\beta$ -arr with siRNA or  $\beta$ -arr encoding plasmids. We identified that  $\beta$ -arr1 silencing and  $\beta$ -arr2 up-regulation were most detrimental for cells and focused on these two conditions for further experiments.

Since  $\beta$ -arr isoforms also play opposing roles on IGF-1R signaling thus we next investigated the effect of  $\beta$ -arr1 silencing and  $\beta$ -arr2 overexpression on receptor signaling [412]. We transfected DFB and BE cells with siRNA against  $\beta$ -arr1 or  $\beta$ -arr2 encoding plasmid followed by serum starvation and IGF-1 stimulation for 0, 2, 5, 10, 30 and 60 minutes. Using WB analysis, we demonstrated that in both  $\beta$ -arr1 silenced and  $\beta$ -arr2 overexpressed cells level of phospho-ERK decreased at 30 and 60 min after stimulation. We further investigated the effect of  $\beta$ -arr modulation on proliferative/survival response of cells to IGF-1. Cells were serum starved after transfection, stimulated or not with IGF-1 for 24 h, followed by cell number analysis. Both  $\beta$ -arr1 downregulation and  $\beta$ -arr2 up-regulation inhibited IGF-1 response by decreased number of cells compared to unstimulated control. These results confirmed that a  $\beta$ -arr2 predominance in melanoma cells inhibits IGF-1 mediated pERK activation as well as an IGF-1 induced proliferative/survival response.


$\beta$ -arr isoforms were demonstrated to induce different patterns of IGF-1R ubiquitination, followed by its degradation or endocytosis. It was also shown that IGF-1R internalization through  $\beta$ -arr1 or  $\beta$ -arr2 has opposing effects on the activation of p53 pathway [272, 282, 412]. Therefore, we next investigated the effect of  $\beta$ -arr modulation on IGF-1R degradation and activity of p53 pathway. Using WB analysis, we identified a decreased rate of IGF-1 induced receptor degradation in both  $\beta$ -arr1 silenced and  $\beta$ -arr2 upregulated cells. Importantly, in both conditions, the level of IGF-1R was decreased at time 0 compared to mock. WB analysis of lysates for p53 expression indicated its upregulation in both conditions. These data demonstrated that  $\beta$ -arr2 predominance downregulates ligand-free IGF-1R, inhibits ligand

induced IGF-1R degradation and activates p53 in melanoma cells.

DTIC is an alkylating agent that induces DNA damage causing cell cycle arrest and inhibition of melanoma cell growth [413]. Thus, we next tested the effect of DTIC as a single agent in BE, DFB and Mel28 melanoma cell lines. Cells were treated for 24 and 48 h with various doses of DTIC in media with or without serum or IGF-1. Using PrestoBlue we demonstrated decreased cell viability and in all cell lines. WB analysis of samples treated with different doses of DTIC for 24 h indicated an increase in the level of p53. DFB cells were more sensitive to DTIC and demonstrated a higher reactivation of p53 than in BE and Mel28 cells. These data confirmed the effect of DTIC on melanoma cells.

Since we demonstrated a negative effect of  $\beta$ -arr2 predominance on cell proliferation, we next tested if this can increase the sensitivity of melanoma cells to DTIC. We compared the effect of single and combined treatments of DTIC with  $\beta$ -arr1 silencing and  $\beta$ -arr2 overexpression. We calculated a predicted response to combined treatment by addition of two single agent sensitivity data. We observed a synergistic effect of combined treatments in all cell lines, indicating that  $\beta$ -arr2 predominance potentiates the DNA damage induced by DTIC in melanoma cells.

The main outcomes of this study are that 1)  $\beta$ -arr2 predominance positively affects p53 activation and negatively affects ligand induced IGF-1R degradation, MAPK activation and proliferation and survival in melanoma cells; 2)  $\beta$ -arr2 predominance synergizes with DTIC treatment in both mtp53 and wtp53 cells.



**Figure 8.** Schematic illustration of the combined effects of dacarbazine with  $\beta$ -arr2 predominance

## 4. DISCUSSION AND FUTURE PROSPECTIVES

The starting point of this thesis was that IGF-1R signaling and p53 pathway share similar regulatory mechanisms: Mdm2 and  $\beta$ arrs. This cross-regulation between two pathways involved in cancer has important implications for anti-cancer therapy and needs to be fully understood. The overall goal of the thesis was to investigate the IGF-1R/ $\beta$ -arr/Mdm2/p53 axis in cancer and to explore the potential use of its components as therapeutic targets.

In the first paper, we investigated the effect of destabilizing IGF-1R/Mdm2/p53 complex using Nutlin-3 and melanoma cells as an experimental model. In the second paper, we focused on the  $\beta$ -arr component of this axis and exploited the possibility to target these components in melanoma. In the third and fourth papers, we tried to exploit different approaches to target the IGF-1R/ $\beta$ -arr/Mdm2/p53 axis. In the third paper, we investigate the effect of co-targeting IGF-1R with MEK pathway, and in the fourth paper we explore the possibility of co-targeting  $\beta$ -arrs with a DNA repair pathway.

Our group has previously described various compounds and antibodies against IGF-1R as IGF-1R biased agonists [305-307]. Anti-IGF-1R antibody CP induces biased pERK that contributes to the survival of CP-treated Ewing's sarcoma cell lines [307]. The antimicrobial peptide LL-37 downregulates the IGF-1R whilst activating biased signaling, that contributes to breast cancer cells invasion and migration [306]. My studies reveal that Nutlin-3 also acts as an IGF-1R biased agonist, activating the MAPK pathway, that contributes to cell migration but not invasion. This study points out a highly specific mechanism of activation of IGF-1R biased signaling by various agonists, resulting in distinct biological outcomes. Characterization of the proteins involved in these pathways that could increase the possibility of more specific anti-cancer targeting is one of the future prospectives of this project. The involvement of  $\beta$ -arr isoforms in this signaling deserves particular attention. It has been demonstrated that  $\beta$ -arr1 mediates CP and LL-37 induced IGF-1R biased signaling [306, 307], however the role of  $\beta$ -arr isoforms in Nutlin-3 mediated ERK activation, as well as the specific role of  $\beta$ -arr2 in any types of IGF-1R signaling has not yet been identified.

In one of the studies we particularly focused on the  $\beta$ -arr components of the IGF-1R/ $\beta$ -arr/Mdm2/p53 axis. We revealed opposing roles of  $\beta$ -arr isoforms in the regulation of IGF-1R at multiple levels –  $\beta$ -arr1 induces while  $\beta$ -arr2 inhibits ligand-dependent IGF-1R degradation;  $\beta$ -arr1 sustains while  $\beta$ -arr2 shortens ERK activation; and  $\beta$ -arr2 activates, while  $\beta$ -arr1 inhibits p53 pathway downstream of IGF-1R activation. However, in the absence of IGF-1 both  $\beta$ -arrs downregulate IGF-1R with  $\beta$ -arr2 exerting a stronger effect. This points out that IGF-1R acts as both a Class A and Class B GPCR depending on the presence of the ligand. It seems that in the absence of the ligand IGF-1R preferentially and transiently recruits  $\beta$ -arr2, while an IGF-1 stimulated IGF-1R stably recruits  $\beta$ -arr1. Another important finding is that the antagonism between the two  $\beta$ -arr isoforms could be used as a target for anti-cancer therapy. We demonstrated that  $\beta$ -arr2 predominance inhibits response to IGF-1, causes cell cycle arrest, and activates p53 pathway resulting in a decrease of cell viability.

Opposing role of  $\beta$ -arrs at different levels of IGF-1R regulation raises the question of the reason for divergence of two isoforms with 78% identity and 88% similarity [414]. Possible



explanations include different affinity of the isoforms for distinct IGF-1R conformations, as well as competition for binding to the receptor and to each other [247, 392, 415, 416]. Another question is whether distinct agonists also induce different conformations of IGF-1R, preferentially recruiting one of the isoforms. In this case biased agonists with preferential recruitment of  $\beta$ -arr2 would be favoured for anti-cancer treatment. Elucidation of the exact mechanism of IGF-1R interaction with  $\beta$ -arr isoforms is one of the future prospectives.

Investigation of the possibility to combine IGF-1R inhibitors with MEK inhibitors indicated that results are critically different depending on the strategy of IGF-1R inhibition. The first strategy of balanced IGF-1R inhibition by siRNA greatly enhanced response to MAPK inhibitors. However, a second strategy using anti-IGF-1R antibody CP was less successful. We revealed that CP not only induces biased IGF-1R signaling, but also is able to bias the IGF-1R towards its ligand even after CP removal. Thus, it was not surprising that CP treatment demonstrated limited effects on MEK1/2 inhibition. As discussed previously, certain RTK antibodies can bind to the receptor's extracellular domain without affecting receptor/ligand interaction [292-296]. Moreover, a specific IR antibody has been described that increases the binding affinity of insulin to the IR thus enhancing the receptors metabolic signaling [301]. This mechanism could be a possible explanation of the specific effect of CP on IGF-1R. The precise mechanism of CP/IGF-1R interaction remains to be investigated. A third strategy of IGF-1R inhibition using Nutlin-3 greatly enhanced the effect of MEK inhibitors. The transient character of Nutlin-3 induced biased signaling is an explanation for the better outcome. Thus, co-targeting MAPK pathway with IGF-1R using Nutlin-3 is the first anti-cancer strategy revealed in this project.

Another study was focused on the possibility to co-target  $\beta$ -arr components of the axis with the p53 pathway. We demonstrated that both  $\beta$ -arr1 inhibition and  $\beta$ -arr2 overexpression increase sensitivity of melanoma cells to DNA damage inducing drug dacarbazine. This study identified the second rational anti-cancer strategy in this project.

In conclusion, two successful strategies of co-targeting IGF-1R/ $\beta$ -arr/Mdm2/p53 axis with different cancer related pathways have been identified in this thesis. One strategy combines MEK inhibition with Nutlin-3 to greatly enhance IGF-1R degradation, transiently activate pERK and induce p53 re-activation. Another strategy is the combination inhibition of  $\beta$ -arr1 or activation of  $\beta$ -arr2 with dacarbazine treatment. This strategy, in contrast, has a moderate effect on IGF-1R degradation, strongly inhibits IGF-1R signaling and has double potential to activate p53 pathway. The synergistic effect of both strategies shows that accurate removal of multiple interconnected pathways of IGF-1R/ $\beta$ -arr/Mdm2/p53 axis can lead to an optimal context dependent anti-cancer effect.

## 5. ACKNOWLEDGEMENTS

To my main supervisor Leonard Girnita for making this thesis possible. Thanks for all of our discussions and for the freedom to have my own interpretations of things. Thanks for always having time to talk about science, but also being a friend and tolerating our half-joke complaints about the electric chair. In a little more than four years you have helped me to understand some fundamental principles of receptor pathology and cancer biology in general. This experience is priceless.

To my co-supervisor Klas Wiman, for giving me an opportunity to work on the p53 project and providing support whenever needed.

To Claire Worrall for introducing me to almost all basics of our research when I first joined the lab. To Iulian Oprea for all the early advice and teachings.

To Caitrin Crudden for making our work together fun, for language corrections, for invasion assays, statistic debates and for random fortune telling skills. Daniela Nedelcu for returning to our lab and teaching me genetic tools. Takashi Shibano, a wizard of immunoprecipitation, for all that I learned from you. To our talented exchange student Benjamin Gebhard for all the help with the drug assays. To external collaborators George Calin and Shin-Ichiro Takahashi. To our clinical team Ada Girnita, Stefan Seregard, Iara Drakensjo, Eric Trocme and to all other present and past members of the group.

To Padraig D'Arcy for the mentorship.

To members of the Wiman lab - Emarndeena, Susanne, Sofia, Vladimir for the materials and advice regarding the p53 part of this project. To other R8:4 corridor-mates - Satendra, Ahmed, Omid, Soniya, Christos etc for making us not feel alone any time in the lab.

To Catharina Svensson for all opportunities I had during my study in Uppsala University.

To my Uppsala-time friends - Anna, Katerina and Pierre for all their influence. To Laetitia for introducing me to GIMP and to the microwaves. Maxwell for books and for the typewriter. Rita for our adventures and dances. To my Vårberg-friends: Sunjay, Fadwa, Yogan, Michalis, Mike etc for all good times together. To my "community friends" - Karen, Natasha Sh, Natasha S, Vladimir, Galya, Nastya and others for our games, dinners and barbeques. To my old friends Tanya, Ayten, Katya, Fidan, Sonya and others for our long friendship.

To my Dad Tofik, to Arzu, to my brothers Ramin and Shamil, sisters-in-law Alma and Aysel and to my wonderful sibsprings Asiya, Timur and Zara. Thanks for your continuous support and happiness that you bring me. To my boyfriend Padraig for an enormous amount of IT-support and for everything else that has become easier and better since we met.

### 3. REFERENCES

1. Jemal, A., et al., *Global cancer statistics*. CA Cancer J Clin, 2011. **61**(2): p. 69-90.
2. Anand, P., et al., *Cancer is a preventable disease that requires major lifestyle changes*. Pharm Res, 2008. **25**(9): p. 2097-116.
3. White, M.C., et al., *Cancer prevention for the next generation*. J Adolesc Health, 2013. **52**(5 Suppl): p. S1-7.
4. Howell, A., et al., *Risk determination and prevention of breast cancer*. Breast Cancer Res, 2014. **16**(5): p. 446.
5. Harvie, M., A. Howell, and D.G. Evans, *Can diet and lifestyle prevent breast cancer: what is the evidence?* Am Soc Clin Oncol Educ Book, 2015: p. e66-73.
6. Hanahan, D. and R.A. Weinberg, *Hallmarks of cancer: the next generation*. Cell, 2011. **144**(5): p. 646-74.
7. Yarden, Y. and A. Ullrich, *Growth factor receptor tyrosine kinases*. Annu Rev Biochem, 1988. **57**: p. 443-78.
8. Witsch, E., M. Sela, and Y. Yarden, *Roles for growth factors in cancer progression*. Physiology (Bethesda), 2010. **25**(2): p. 85-101.
9. Goustin, A.S., et al., *Growth factors and cancer*. Cancer Res, 1986. **46**(3): p. 1015-29.
10. Perona, R., *Cell signalling: growth factors and tyrosine kinase receptors*. Clin Transl Oncol, 2006. **8**(2): p. 77-82.
11. Sun, W. and J. Yang, *Functional mechanisms for human tumor suppressors*. J Cancer, 2010. **1**: p. 136-40.
12. Kruiswijk, F., C.F. Labuschagne, and K.H. Vousden, *p53 in survival, death and metabolic health: a lifeguard with a licence to kill*. Nat Rev Mol Cell Biol, 2015. **16**(7): p. 393-405.
13. Harbour, J.W. and D.C. Dean, *Rb function in cell-cycle regulation and apoptosis*. Nat Cell Biol, 2000. **2**(4): p. E65-7.
14. Sherr, C.J. and F. McCormick, *The RB and p53 pathways in cancer*. Cancer Cell, 2002. **2**(2): p. 103-12.
15. Deshpande, A., P. Sicinski, and P.W. Hinds, *Cyclins and cdks in development and cancer: a perspective*. Oncogene, 2005. **24**(17): p. 2909-15.
16. Tagoug, I., A. Sauty De Chalon, and C. Dumontet, *Inhibition of IGF-1 signalling enhances the apoptotic effect of AS602868, an IKK2 inhibitor, in multiple myeloma cell lines*. PLoS One, 2011. **6**(7): p. e22641.
17. Hoffman, B. and D.A. Liebermann, *Apoptotic signaling by c-MYC*. Oncogene, 2008. **27**(50): p. 6462-72.
18. Igney, F.H. and P.H. Krammer, *Death and anti-death: tumour resistance to apoptosis*. Nat Rev Cancer, 2002. **2**(4): p. 277-88.
19. White, E. and R.S. DiPaola, *The double-edged sword of autophagy modulation in cancer*. Clin Cancer Res, 2009. **15**(17): p. 5308-16.
20. Grivennikov, S.I., F.R. Greten, and M. Karin, *Immunity, inflammation, and cancer*. Cell, 2010. **140**(6): p. 883-99.
21. Blasco, M.A., *Telomeres and human disease: ageing, cancer and beyond*. Nat Rev Genet, 2005. **6**(8): p. 611-22.
22. Shay, J.W. and W.E. Wright, *Role of telomeres and telomerase in cancer*. Semin Cancer Biol, 2011. **21**(6): p. 349-53.
23. Shamas, M.A., *Telomeres, lifestyle, cancer, and aging*. Curr Opin Clin Nutr Metab Care, 2011. **14**(1): p. 28-34.

24. Artandi, S.E. and R.A. DePinho, *Telomeres and telomerase in cancer*. Carcinogenesis, 2010. **31**(1): p. 9-18.
25. Artandi, S.E. and R.A. DePinho, *Mice without telomerase: what can they teach us about human cancer?* Nat Med, 2000. **6**(8): p. 852-5.
26. Papetti, M. and I.M. Herman, *Mechanisms of normal and tumor-derived angiogenesis*. Am J Physiol Cell Physiol, 2002. **282**(5): p. C947-70.
27. Carmeliet, P., *VEGF as a key mediator of angiogenesis in cancer*. Oncology, 2005. **69 Suppl 3**: p. 4-10.
28. Turner, N. and R. Grose, *Fibroblast growth factor signalling: from development to cancer*. Nat Rev Cancer, 2010. **10**(2): p. 116-29.
29. Kazerounian, S., K.O. Yee, and J. Lawler, *Thrombospondins in cancer*. Cell Mol Life Sci, 2008. **65**(5): p. 700-12.
30. Murdoch, C., et al., *The role of myeloid cells in the promotion of tumour angiogenesis*. Nat Rev Cancer, 2008. **8**(8): p. 618-31.
31. Zmsteg, A. and G. Christofori, *Corrupt policemen: inflammatory cells promote tumor angiogenesis*. Curr Opin Oncol, 2009. **21**(1): p. 60-70.
32. Geiger, B. and K.M. Yamada, *Molecular architecture and function of matrix adhesions*. Cold Spring Harb Perspect Biol, 2011. **3**(5).
33. Berx, G. and F. van Roy, *Involvement of members of the cadherin superfamily in cancer*. Cold Spring Harb Perspect Biol, 2009. **1**(6): p. a003129.
34. Yang, J. and R.A. Weinberg, *Epithelial-mesenchymal transition: at the crossroads of development and tumor metastasis*. Dev Cell, 2008. **14**(6): p. 818-29.
35. Joyce, J.A. and J.W. Pollard, *Microenvironmental regulation of metastasis*. Nat Rev Cancer, 2009. **9**(4): p. 239-52.
36. Yang, S.Y., et al., *Growth Factors and their receptors in cancer metastases*. Front Biosci (Landmark Ed), 2011. **16**: p. 531-8.
37. Huttemann, M., et al., *Regulation of mitochondrial oxidative phosphorylation through cell signaling*. Biochim Biophys Acta, 2007. **1773**(12): p. 1701-20.
38. Warburg, O., *On respiratory impairment in cancer cells*. Science, 1956. **124**(3215): p. 269-70.
39. Vander Heiden, M.G., L.C. Cantley, and C.B. Thompson, *Understanding the Warburg effect: the metabolic requirements of cell proliferation*. Science, 2009. **324**(5930): p. 1029-33.
40. Feron, O., *Pyruvate into lactate and back: from the Warburg effect to symbiotic energy fuel exchange in cancer cells*. Radiother Oncol, 2009. **92**(3): p. 329-33.
41. DeBerardinis, R.J., et al., *The biology of cancer: metabolic reprogramming fuels cell growth and proliferation*. Cell Metab, 2008. **7**(1): p. 11-20.
42. Jones, R.G. and C.B. Thompson, *Tumor suppressors and cell metabolism: a recipe for cancer growth*. Genes Dev, 2009. **23**(5): p. 537-48.
43. Diefenbach, A. and D.H. Raulet, *The innate immune response to tumors and its role in the induction of T-cell immunity*. Immunol Rev, 2002. **188**: p. 9-21.
44. Gajewski, T.F., H. Schreiber, and Y.X. Fu, *Innate and adaptive immune cells in the tumor microenvironment*. Nat Immunol, 2013. **14**(10): p. 1014-22.
45. Yang, L., Y. Pang, and H.L. Moses, *TGF-beta and immune cells: an important regulatory axis in the tumor microenvironment and progression*. Trends Immunol, 2010. **31**(6): p. 220-7.
46. Shields, J.D., et al., *Induction of lymphoidlike stroma and immune escape by tumors that express the chemokine CCL21*. Science, 2010. **328**(5979): p. 749-52.
47. Mougiakakos, D., et al., *Regulatory T cells in cancer*. Adv Cancer Res, 2010. **107**: p. 57-117.

48. Ostrand-Rosenberg, S. and P. Sinha, *Myeloid-derived suppressor cells: linking inflammation and cancer*. J Immunol, 2009. **182**(8): p. 4499-506.
49. Hanahan, D. and R.A. Weinberg, *The hallmarks of cancer*. Cell, 2000. **100**(1): p. 57-70.
50. Negrini, S., V.G. Gorgoulis, and T.D. Halazonetis, *Genomic instability--an evolving hallmark of cancer*. Nat Rev Mol Cell Biol, 2010. **11**(3): p. 220-8.
51. Haupt, S., D. Raghu, and Y. Haupt, *Mutant p53 Drives Cancer by Subverting Multiple Tumor Suppression Pathways*. Front Oncol, 2016. **6**: p. 12.
52. DeNardo, D.G., P. Andreu, and L.M. Coussens, *Interactions between lymphocytes and myeloid cells regulate pro- versus anti-tumor immunity*. Cancer Metastasis Rev, 2010. **29**(2): p. 309-16.
53. Qian, B.Z. and J.W. Pollard, *Macrophage diversity enhances tumor progression and metastasis*. Cell, 2010. **141**(1): p. 39-51.
54. Whiteside, T.L., *The tumor microenvironment and its role in promoting tumor growth*. Oncogene, 2008. **27**(45): p. 5904-12.
55. Quail, D.F. and J.A. Joyce, *Microenvironmental regulation of tumor progression and metastasis*. Nat Med, 2013. **19**(11): p. 1423-37.
56. Luo, J., N.L. Solimini, and S.J. Elledge, *Principles of cancer therapy: oncogene and non-oncogene addiction*. Cell, 2009. **136**(5): p. 823-37.
57. Huang, J.S. and S.S. Huang, *Role of growth factors in oncogenesis: growth factor-proto-oncogene pathways of mitogenesis*. Ciba Found Symp, 1985. **116**: p. 46-65.
58. Mosesson, Y. and Y. Yarden, *Oncogenic growth factor receptors: implications for signal transduction therapy*. Semin Cancer Biol, 2004. **14**(4): p. 262-70.
59. Giancotti, F.G., *Deregulation of cell signaling in cancer*. FEBS Lett, 2014. **588**(16): p. 2558-70.
60. Liu, Y., et al., *Targeting tumor suppressor genes for cancer therapy*. Bioessays, 2015. **37**(12): p. 1277-86.
61. Lee, E.Y. and W.J. Muller, *Oncogenes and tumor suppressor genes*. Cold Spring Harb Perspect Biol, 2010. **2**(10): p. a003236.
62. Prenzel, N., et al., *The epidermal growth factor receptor family as a central element for cellular signal transduction and diversification*. Endocr Relat Cancer, 2001. **8**(1): p. 11-31.
63. Lof, E.B., *Growth factor receptor signalling: location, location, location*. Trends Cell Biol, 2000. **10**(8): p. 343-8.
64. Van Obberghen, E., *Signalling through the insulin receptor and the insulin-like growth factor-I receptor*. Diabetologia, 1994. **37 Suppl 2**: p. S125-34.
65. Coleman, S.J., et al., *The ins and outs of fibroblast growth factor receptor signalling*. Clin Sci (Lond), 2014. **127**(4): p. 217-31.
66. Sporn, M.B. and A.B. Roberts, *Autocrine growth factors and cancer*. Nature, 1985. **313**(6005): p. 745-7.
67. Nicolson, G.L., *Paracrine/autocrine growth mechanisms in tumor metastasis*. Oncol Res, 1992. **4**(10): p. 389-99.
68. Buchholz, T.A., et al., *Tumor suppressor genes and breast cancer*. Radiat Oncol Investig, 1999. **7**(2): p. 55-65.
69. Berger, A.H., A.G. Knudson, and P.P. Pandolfi, *A continuum model for tumour suppression*. Nature, 2011. **476**(7359): p. 163-9.
70. Aaronson, S.A., *Growth factors and cancer*. Science, 1991. **254**(5035): p. 1146-53.
71. Schlessinger, J., *Cell signaling by receptor tyrosine kinases*. Cell, 2000. **103**(2): p. 211-25.

72. Robinson, D.R., Y.M. Wu, and S.F. Lin, *The protein tyrosine kinase family of the human genome*. *Oncogene*, 2000. **19**(49): p. 5548-57.
73. Hubbard, S.R., M. Mohammadi, and J. Schlessinger, *Autoregulatory mechanisms in protein-tyrosine kinases*. *J Biol Chem*, 1998. **273**(20): p. 11987-90.
74. Ullrich, A. and J. Schlessinger, *Signal transduction by receptors with tyrosine kinase activity*. *Cell*, 1990. **61**(2): p. 203-12.
75. Schlessinger, J. and M.A. Lemmon, *SH2 and PTB domains in tyrosine kinase signaling*. *Sci STKE*, 2003. **2003**(191): p. RE12.
76. Pawson, T., *Specificity in signal transduction: from phosphotyrosine-SH2 domain interactions to complex cellular systems*. *Cell*, 2004. **116**(2): p. 191-203.
77. Noselli, S. and N. Perrimon, *Signal transduction. Are there close encounters between signaling pathways?* *Science*, 2000. **290**(5489): p. 68-9.
78. Aaronson, S.A., et al., *Growth factors and malignant transformation*. *Adv Exp Med Biol*, 1993. **348**: p. 7-22.
79. Wieduwilt, M.J. and M.M. Moasser, *The epidermal growth factor receptor family: biology driving targeted therapeutics*. *Cell Mol Life Sci*, 2008. **65**(10): p. 1566-84.
80. Sasaki, T., K. Hiroki, and Y. Yamashita, *The role of epidermal growth factor receptor in cancer metastasis and microenvironment*. *Biomed Res Int*, 2013. **2013**: p. 546318.
81. Simons, M., E. Gordon, and L. Claesson-Welsh, *Mechanisms and regulation of endothelial VEGF receptor signalling*. *Nat Rev Mol Cell Biol*, 2016. **17**(10): p. 611-25.
82. Olsson, A.K., et al., *VEGF receptor signalling - in control of vascular function*. *Nat Rev Mol Cell Biol*, 2006. **7**(5): p. 359-71.
83. Shibuya, M., *Vascular Endothelial Growth Factor (VEGF) and Its Receptor (VEGFR) Signaling in Angiogenesis: A Crucial Target for Anti- and Pro-Angiogenic Therapies*. *Genes Cancer*, 2011. **2**(12): p. 1097-105.
84. Rapisarda, A. and G. Melillo, *Role of the VEGF/VEGFR axis in cancer biology and therapy*. *Adv Cancer Res*, 2012. **114**: p. 237-67.
85. Andrae, J., R. Gallini, and C. Betsholtz, *Role of platelet-derived growth factors in physiology and medicine*. *Genes Dev*, 2008. **22**(10): p. 1276-312.
86. Yu, J., C. Ustach, and H.R. Kim, *Platelet-derived growth factor signaling and human cancer*. *J Biochem Mol Biol*, 2003. **36**(1): p. 49-59.
87. Demoulin, J.B. and A. Essaghir, *PDGF receptor signaling networks in normal and cancer cells*. *Cytokine Growth Factor Rev*, 2014. **25**(3): p. 273-83.
88. Girnita, L., et al., *Something old, something new and something borrowed: emerging paradigm of insulin-like growth factor type 1 receptor (IGF-1R) signaling regulation*. *Cell Mol Life Sci*, 2014. **71**(13): p. 2403-27.
89. Li, R., A. Pourpak, and S.W. Morris, *Inhibition of the insulin-like growth factor-1 receptor (IGF1R) tyrosine kinase as a novel cancer therapy approach*. *J Med Chem*, 2009. **52**(16): p. 4981-5004.
90. Romero, C.J., et al., *Insulin-like growth factor 1 mediates negative feedback to somatotroph GH expression via POU1F1/CREB binding protein interactions*. *Mol Cell Biol*, 2012. **32**(21): p. 4258-69.
91. Arany, E., et al., *Differential cellular synthesis of insulin-like growth factor binding protein-1 (IGFBP-1) and IGFBP-3 within human liver*. *J Clin Endocrinol Metab*, 1994. **79**(6): p. 1871-6.
92. Olivecrona, H., et al., *Acute and short-term effects of growth hormone on insulin-like growth factors and their binding proteins: serum levels and hepatic messenger ribonucleic acid responses in humans*. *J Clin Endocrinol Metab*, 1999. **84**(2): p. 553-60.

93. Adams, T.E., et al., *Structure and function of the type I insulin-like growth factor receptor*. Cell Mol Life Sci, 2000. **57**(7): p. 1050-93.
94. Carlberg, M., et al., *Mevalonic acid is limiting for N-linked glycosylation and translocation of the insulin-like growth factor-I receptor to the cell surface. Evidence for a new link between 3-hydroxy-3-methylglutaryl-coenzyme A reductase and cell growth*. J Biol Chem, 1996. **271**(29): p. 17453-62.
95. Sepp-Lorenzino, L., *Structure and function of the insulin-like growth factor I receptor*. Breast Cancer Res Treat, 1998. **47**(3): p. 235-53.
96. Ullrich, A., et al., *Insulin-like growth factor I receptor primary structure: comparison with insulin receptor suggests structural determinants that define functional specificity*. EMBO J, 1986. **5**(10): p. 2503-12.
97. Favelyukis, S., et al., *Structure and autoregulation of the insulin-like growth factor I receptor kinase*. Nat Struct Biol, 2001. **8**(12): p. 1058-63.
98. Butler, A.A., et al., *Insulin-like growth factor-I receptor signal transduction: at the interface between physiology and cell biology*. Comp Biochem Physiol B Biochem Mol Biol, 1998. **121**(1): p. 19-26.
99. Werner, H. and D. Le Roith, *New concepts in regulation and function of the insulin-like growth factors: implications for understanding normal growth and neoplasia*. Cell Mol Life Sci, 2000. **57**(6): p. 932-42.
100. Wick, M.J., et al., *Mechanism of phosphorylation of protein kinase B/Akt by a constitutively active 3-phosphoinositide-dependent protein kinase-1*. J Biol Chem, 2000. **275**(51): p. 40400-6.
101. Balendran, A., et al., *Evidence that 3-phosphoinositide-dependent protein kinase-1 mediates phosphorylation of p70 S6 kinase in vivo at Thr-412 as well as Thr-252*. J Biol Chem, 1999. **274**(52): p. 37400-6.
102. Fukushima, T., et al., *Phosphatidylinositol 3-kinase (PI3K) activity bound to insulin-like growth factor-I (IGF-I) receptor, which is continuously sustained by IGF-I stimulation, is required for IGF-I-induced cell proliferation*. J Biol Chem, 2012. **287**(35): p. 29713-21.
103. Chong, H., H.G. Vikis, and K.L. Guan, *Mechanisms of regulating the Raf kinase family*. Cell Signal, 2003. **15**(5): p. 463-9.
104. Vetter, I.R. and A. Wittinghofer, *The guanine nucleotide-binding switch in three dimensions*. Science, 2001. **294**(5545): p. 1299-304.
105. Bonni, A., et al., *Cell survival promoted by the Ras-MAPK signaling pathway by transcription-dependent and -independent mechanisms*. Science, 1999. **286**(5443): p. 1358-62.
106. Zhang, W. and H.T. Liu, *MAPK signal pathways in the regulation of cell proliferation in mammalian cells*. Cell Res, 2002. **12**(1): p. 9-18.
107. Yoon, S. and R. Seger, *The extracellular signal-regulated kinase: multiple substrates regulate diverse cellular functions*. Growth Factors, 2006. **24**(1): p. 21-44.
108. Peruzzi, F., et al., *Multiple signaling pathways of the insulin-like growth factor I receptor in protection from apoptosis*. Mol Cell Biol, 1999. **19**(10): p. 7203-15.
109. Vincent, A.M. and E.L. Feldman, *Control of cell survival by IGF signaling pathways*. Growth Horm IGF Res, 2002. **12**(4): p. 193-7.
110. Mauro, L., et al., *Role of the IGF-I receptor in the regulation of cell-cell adhesion: implications in cancer development and progression*. J Cell Physiol, 2003. **194**(2): p. 108-16.
111. Khandwala, H.M., et al., *The effects of insulin-like growth factors on tumorigenesis and neoplastic growth*. Endocr Rev, 2000. **21**(3): p. 215-44.

112. Liu, J.P., et al., *Mice carrying null mutations of the genes encoding insulin-like growth factor I (Igf-I) and type I IGF receptor (Igf1r)*. Cell, 1993. **75**(1): p. 59-72.
113. Pollak, M.N., et al., *Presence of somatomedin receptors on primary human breast and colon carcinomas*. Cancer Lett, 1987. **38**(1-2): p. 223-30.
114. Baserga, R., *The insulin-like growth factor I receptor: a key to tumor growth?* Cancer Res, 1995. **55**(2): p. 249-52.
115. Larsson, O., A. Girnita, and L. Girnita, *Role of insulin-like growth factor I receptor signalling in cancer*. Br J Cancer, 2007. **96 Suppl**: p. R2-6.
116. Chan, J.M., et al., *Insulin-like growth factor-I (IGF-I) and IGF binding protein-3 as predictors of advanced-stage prostate cancer*. J Natl Cancer Inst, 2002. **94**(14): p. 1099-106.
117. Dunn, S.E., et al., *A dominant negative mutant of the insulin-like growth factor-I receptor inhibits the adhesion, invasion, and metastasis of breast cancer*. Cancer Res, 1998. **58**(15): p. 3353-61.
118. Hankinson, S.E., et al., *Circulating concentrations of insulin-like growth factor-I and risk of breast cancer*. Lancet, 1998. **351**(9113): p. 1393-6.
119. Pollak, M.N., E.S. Schernhammer, and S.E. Hankinson, *Insulin-like growth factors and neoplasia*. Nat Rev Cancer, 2004. **4**(7): p. 505-18.
120. Brahmkhatri, V.P., C. Prasanna, and H.S. Atreya, *Insulin-like growth factor system in cancer: novel targeted therapies*. Biomed Res Int, 2015. **2015**: p. 538019.
121. Grimberg, A., *Mechanisms by which IGF-I may promote cancer*. Cancer Biol Ther, 2003. **2**(6): p. 630-5.
122. LeRoith, D. and C.T. Roberts, Jr., *The insulin-like growth factor system and cancer*. Cancer Lett, 2003. **195**(2): p. 127-37.
123. Lee, A.V., et al., *Enhancement of insulin-like growth factor signaling in human breast cancer: estrogen regulation of insulin receptor substrate-1 expression in vitro and in vivo*. Mol Endocrinol, 1999. **13**(5): p. 787-96.
124. Vorwerk, P., et al., *Expression of components of the IGF signalling system in childhood acute lymphoblastic leukaemia*. Mol Pathol, 2002. **55**(1): p. 40-5.
125. McGuire, W.L., Jr., et al., *Regulation of insulin-like growth factor-binding protein (IGFBP) expression by breast cancer cells: use of IGFBP-1 as an inhibitor of insulin-like growth factor action*. J Natl Cancer Inst, 1992. **84**(17): p. 1336-41.
126. Karas, M., et al., *Membrane-associated insulin-like growth factor-binding protein-3 inhibits insulin-like growth factor-I-induced insulin-like growth factor-I receptor signaling in ishikawa endometrial cancer cells*. J Biol Chem, 1997. **272**(26): p. 16514-20.
127. Yu, H. and T. Rohan, *Role of the insulin-like growth factor family in cancer development and progression*. J Natl Cancer Inst, 2000. **92**(18): p. 1472-89.
128. Roberts, C.T., Jr., *IGF-I and prostate cancer*. Novartis Found Symp, 2004. **262**: p. 193-9; discussion 199-204, 265-8.
129. Chott, A., et al., *Tyrosine kinases expressed in vivo by human prostate cancer bone marrow metastases and loss of the type I insulin-like growth factor receptor*. Am J Pathol, 1999. **155**(4): p. 1271-9.
130. Mancarella, C., et al., *Insulin-like growth factor I receptor affects the survival of primary prostate cancer patients depending on TMPRSS2-ERG status*. BMC Cancer, 2017. **17**(1): p. 367.
131. Tennant, M.K., et al., *Protein and messenger ribonucleic acid (mRNA) for the type I insulin-like growth factor (IGF) receptor is decreased and IGF-II mRNA is increased in human prostate carcinoma compared to benign prostate epithelium*. J Clin Endocrinol Metab, 1996. **81**(10): p. 3774-82.



132. Baserga, R. and R. Rubin, *Cell cycle and growth control*. Crit Rev Eukaryot Gene Expr, 1993. **3**(1): p. 47-61.
133. Kanter-Lewensohn, L., et al., *Expression of insulin-like growth factor-1 receptor (IGF-1R) and p27Kip1 in melanocytic tumors: a potential regulatory role of IGF-1 pathway in distribution of p27Kip1 between different cyclins*. Growth Factors, 2000. **17**(3): p. 193-202.
134. Reiss, K., et al., *Insulin-like growth factor-1 receptor and its ligand regulate the reentry of adult ventricular myocytes into the cell cycle*. Exp Cell Res, 1997. **235**(1): p. 198-209.
135. Baserga, R. and A. Morrión, *Differentiation and malignant transformation: two roads diverged in a wood*. J Cell Biochem, 1999. **Suppl 32-33**: p. 68-75.
136. Sell, C., et al., *Effect of a null mutation of the insulin-like growth factor I receptor gene on growth and transformation of mouse embryo fibroblasts*. Mol Cell Biol, 1994. **14**(6): p. 3604-12.
137. Valentinis, B., et al., *Insulin-like growth factor I receptor signaling in transformation by src oncogenes*. Mol Cell Biol, 1997. **17**(7): p. 3744-54.
138. Liu, J.L., et al., *The constitutively active mutant Galpha13 transforms mouse fibroblast cells deficient in insulin-like growth factor-I receptor*. J Biol Chem, 1997. **272**(47): p. 29438-41.
139. DeAngelis, T., et al., *Transformation by the simian virus 40 T antigen is regulated by IGF-I receptor and IRS-1 signaling*. Oncogene, 2006. **25**(1): p. 32-42.
140. Gajewski, T.F. and C.B. Thompson, *Apoptosis meets signal transduction: elimination of a BAD influence*. Cell, 1996. **87**(4): p. 589-92.
141. Dudek, H., et al., *Regulation of neuronal survival by the serine-threonine protein kinase Akt*. Science, 1997. **275**(5300): p. 661-5.
142. Kauffmann-Zeh, A., et al., *Suppression of c-Myc-induced apoptosis by Ras signalling through PI(3)K and PKB*. Nature, 1997. **385**(6616): p. 544-8.
143. O'Connor, R., et al., *Identification of domains of the insulin-like growth factor I receptor that are required for protection from apoptosis*. Mol Cell Biol, 1997. **17**(1): p. 427-35.
144. Hongo, A., et al., *Mutational analysis of the mitogenic and transforming activities of the insulin-like growth factor I receptor*. Oncogene, 1996. **12**(6): p. 1231-8.
145. Cheng, H.L., et al., *GTPases and phosphatidylinositol 3-kinase are critical for insulin-like growth factor-I-mediated Schwann cell motility*. J Biol Chem, 2000. **275**(35): p. 27197-204.
146. Cheng, H.L., et al., *IGF-I promotes Schwann cell motility and survival via activation of Akt*. Mol Cell Endocrinol, 2000. **170**(1-2): p. 211-5.
147. Buckley, D.A., et al., *Regulation of insulin-like growth factor type I (IGF-I) receptor kinase activity by protein tyrosine phosphatase 1B (PTP-1B) and enhanced IGF-I-mediated suppression of apoptosis and motility in PTP-1B-deficient fibroblasts*. Mol Cell Biol, 2002. **22**(7): p. 1998-2010.
148. Zhang, D. and P. Brodt, *Type I insulin-like growth factor regulates MT1-MMP synthesis and tumor invasion via PI 3-kinase/Akt signaling*. Oncogene, 2003. **22**(7): p. 974-82.
149. Girnita, A., et al., *The insulin-like growth factor-I receptor inhibitor picropodophyllin causes tumor regression and attenuates mechanisms involved in invasion of uveal melanoma cells*. Clin Cancer Res, 2006. **12**(4): p. 1383-91.
150. Casamassima, A. and E. Rozengurt, *Insulin-like growth factor I stimulates tyrosine phosphorylation of p130(Cas), focal adhesion kinase, and paxillin. Role of*

- phosphatidylinositol 3'-kinase and formation of a p130(Cas).Crk complex.* J Biol Chem, 1998. **273**(40): p. 26149-56.
151. Beitner-Johnson, D. and D. LeRoith, *Insulin-like growth factor-I stimulates tyrosine phosphorylation of endogenous c-Crk.* J Biol Chem, 1995. **270**(10): p. 5187-90.
  152. Sekharam, M., et al., *Insulin-like growth factor 1 receptor activates c-SRC and modifies transformation and motility of colon cancer in vitro.* Anticancer Res, 2003. **23**(2B): p. 1517-24.
  153. Felding-Habermann, B., *Integrin adhesion receptors in tumor metastasis.* Clin Exp Metastasis, 2003. **20**(3): p. 203-13.
  154. Zhang, D., et al., *Dual regulation of MMP-2 expression by the type 1 insulin-like growth factor receptor: the phosphatidylinositol 3-kinase/Akt and Raf/ERK pathways transmit opposing signals.* J Biol Chem, 2004. **279**(19): p. 19683-90.
  155. Feldser, D., et al., *Reciprocal positive regulation of hypoxia-inducible factor 1alpha and insulin-like growth factor 2.* Cancer Res, 1999. **59**(16): p. 3915-8.
  156. Zelzer, E., et al., *Insulin induces transcription of target genes through the hypoxia-inducible factor HIF-1alpha/ARNT.* EMBO J, 1998. **17**(17): p. 5085-94.
  157. Clarke, K., et al., *Mutant epidermal growth factor receptor enhances induction of vascular endothelial growth factor by hypoxia and insulin-like growth factor-1 via a PI3 kinase dependent pathway.* Br J Cancer, 2001. **84**(10): p. 1322-9.
  158. Dunn, S.E., et al., *Up-regulation of urokinase-type plasminogen activator by insulin-like growth factor-1 depends upon phosphatidylinositol-3 kinase and mitogen-activated protein kinase kinase.* Cancer Res, 2001. **61**(4): p. 1367-74.
  159. Lopez, T. and D. Hanahan, *Elevated levels of IGF-1 receptor convey invasive and metastatic capability in a mouse model of pancreatic islet tumorigenesis.* Cancer Cell, 2002. **1**(4): p. 339-53.
  160. Samani, A.A., et al., *The role of the IGF system in cancer growth and metastasis: overview and recent insights.* Endocr Rev, 2007. **28**(1): p. 20-47.
  161. Fredriksson, R., et al., *The G-protein-coupled receptors in the human genome form five main families. Phylogenetic analysis, paralogon groups, and fingerprints.* Mol Pharmacol, 2003. **63**(6): p. 1256-72.
  162. Hamm, H.E., *How activated receptors couple to G proteins.* Proc Natl Acad Sci U S A, 2001. **98**(9): p. 4819-21.
  163. Lefkowitz, R.J., *Historical review: a brief history and personal retrospective of seven-transmembrane receptors.* Trends Pharmacol Sci, 2004. **25**(8): p. 413-22.
  164. Bos, J.L., H. Rehmann, and A. Wittinghofer, *GEFs and GAPs: critical elements in the control of small G proteins.* Cell, 2007. **129**(5): p. 865-77.
  165. Oldham, W.M. and H.E. Hamm, *Heterotrimeric G protein activation by G-protein-coupled receptors.* Nat Rev Mol Cell Biol, 2008. **9**(1): p. 60-71.
  166. Penela, P., C. Ribas, and F. Mayor, Jr., *Mechanisms of regulation of the expression and function of G protein-coupled receptor kinases.* Cell Signal, 2003. **15**(11): p. 973-81.
  167. Perry, S.J., et al., *Targeting of cyclic AMP degradation to beta 2-adrenergic receptors by beta-arrestins.* Science, 2002. **298**(5594): p. 834-6.
  168. Klein, S., H. Reuveni, and A. Levitzki, *Signal transduction by a nondissociable heterotrimeric yeast G protein.* Proc Natl Acad Sci U S A, 2000. **97**(7): p. 3219-23.
  169. Bunemann, M., M. Frank, and M.J. Lohse, *Gi protein activation in intact cells involves subunit rearrangement rather than dissociation.* Proc Natl Acad Sci U S A, 2003. **100**(26): p. 16077-82.

170. Gales, C., et al., *Probing the activation-promoted structural rearrangements in preassembled receptor-G protein complexes*. Nat Struct Mol Biol, 2006. **13**(9): p. 778-86.
171. Rebois, R.V., D.R. Warner, and N.S. Basi, *Does subunit dissociation necessarily accompany the activation of all heterotrimeric G proteins?* Cell Signal, 1997. **9**(2): p. 141-51.
172. Cahill, T.J., 3rd, et al., *Distinct conformations of GPCR-beta-arrestin complexes mediate desensitization, signaling, and endocytosis*. Proc Natl Acad Sci U S A, 2017. **114**(10): p. 2562-2567.
173. Thomsen, A.R., et al., *GPCR-G Protein-beta-Arrestin Super-Complex Mediates Sustained G Protein Signaling*. Cell, 2016. **166**(4): p. 907-19.
174. Lohse, M.J., et al., *Kinetics of G-protein-coupled receptor signals in intact cells*. Br J Pharmacol, 2008. **153 Suppl 1**: p. S125-32.
175. Smrcka, A.V., *G protein betagamma subunits: central mediators of G protein-coupled receptor signaling*. Cell Mol Life Sci, 2008. **65**(14): p. 2191-214.
176. Maudsley, S., B. Martin, and L.M. Luttrell, *The origins of diversity and specificity in g protein-coupled receptor signaling*. J Pharmacol Exp Ther, 2005. **314**(2): p. 485-94.
177. Simon, M.I., M.P. Strathmann, and N. Gautam, *Diversity of G proteins in signal transduction*. Science, 1991. **252**(5007): p. 802-8.
178. Guo, Y., et al., *Predicting G-protein coupled receptors-G-protein coupling specificity based on autocross-covariance transform*. Proteins, 2006. **65**(1): p. 55-60.
179. Sgourakis, N.G., et al., *A method for the prediction of GPCRs coupling specificity to G-proteins using refined profile Hidden Markov Models*. BMC Bioinformatics, 2005. **6**: p. 104.
180. Cabrera-Vera, T.M., et al., *Insights into G protein structure, function, and regulation*. Endocr Rev, 2003. **24**(6): p. 765-81.
181. Pierce, K.L., R.T. Premont, and R.J. Lefkowitz, *Seven-transmembrane receptors*. Nat Rev Mol Cell Biol, 2002. **3**(9): p. 639-50.
182. Rasmussen, S.G., et al., *Crystal structure of the beta2 adrenergic receptor-Gs protein complex*. Nature, 2011. **477**(7366): p. 549-55.
183. Albarran-Juarez, J., et al., *Modulation of alpha2-adrenoceptor functions by heterotrimeric Galphai protein isoforms*. J Pharmacol Exp Ther, 2009. **331**(1): p. 35-44.
184. Melsom, C.B., et al., *Non-classical regulation of beta1- and beta 2-adrenoceptor-mediated inotropic responses in rat heart ventricle by the G protein Gi*. Naunyn Schmiedebergs Arch Pharmacol, 2014. **387**(12): p. 1177-86.
185. Macrez-Lepretre, N., et al., *Distinct functions of Gq and G11 proteins in coupling alpha1-adrenoreceptors to Ca2+ release and Ca2+ entry in rat portal vein myocytes*. J Biol Chem, 1997. **272**(8): p. 5261-8.
186. Guo, D.F., et al., *The angiotensin II type 1 receptor and receptor-associated proteins*. Cell Res, 2001. **11**(3): p. 165-80.
187. McCoy, K.L., S.F. Traynelis, and J.R. Hepler, *PAR1 and PAR2 couple to overlapping and distinct sets of G proteins and linked signaling pathways to differentially regulate cell physiology*. Mol Pharmacol, 2010. **77**(6): p. 1005-15.
188. Siehler, S., *Regulation of RhoGEF proteins by G12/13-coupled receptors*. Br J Pharmacol, 2009. **158**(1): p. 41-9.
189. Eason, M.G., et al., *Simultaneous coupling of alpha 2-adrenergic receptors to two G-proteins with opposing effects. Subtype-selective coupling of alpha 2C10, alpha 2C4, and alpha 2C2 adrenergic receptors to Gi and Gs*. J Biol Chem, 1992. **267**(22): p. 15795-801.

190. Rajagopal, S., K. Rajagopal, and R.J. Lefkowitz, *Teaching old receptors new tricks: biasing seven-transmembrane receptors*. Nat Rev Drug Discov, 2010. **9**(5): p. 373-86.
191. Clapham, D.E. and E.J. Neer, *G protein beta gamma subunits*. Annu Rev Pharmacol Toxicol, 1997. **37**: p. 167-203.
192. Pitcher, J.A., et al., *Pleckstrin homology domain-mediated membrane association and activation of the beta-adrenergic receptor kinase requires coordinate interaction with G beta gamma subunits and lipid*. J Biol Chem, 1995. **270**(20): p. 11707-10.
193. Daaka, Y., et al., *Receptor and G betagamma isoform-specific interactions with G protein-coupled receptor kinases*. Proc Natl Acad Sci U S A, 1997. **94**(6): p. 2180-5.
194. Pitcher, J.A., et al., *Feedback inhibition of G protein-coupled receptor kinase 2 (GRK2) activity by extracellular signal-regulated kinases*. J Biol Chem, 1999. **274**(49): p. 34531-4.
195. Gurevich, E.V., et al., *G protein-coupled receptor kinases: more than just kinases and not only for GPCRs*. Pharmacol Ther, 2012. **133**(1): p. 40-69.
196. Ribas, C., et al., *The G protein-coupled receptor kinase (GRK) interactome: role of GRKs in GPCR regulation and signaling*. Biochim Biophys Acta, 2007. **1768**(4): p. 913-22.
197. Bunemann, M. and M.M. Hosey, *G-protein coupled receptor kinases as modulators of G-protein signalling*. J Physiol, 1999. **517** ( Pt 1): p. 5-23.
198. Jaber, M., et al., *Essential role of beta-adrenergic receptor kinase 1 in cardiac development and function*. Proc Natl Acad Sci U S A, 1996. **93**(23): p. 12974-9.
199. Kohout, T.A. and R.J. Lefkowitz, *Regulation of G protein-coupled receptor kinases and arrestins during receptor desensitization*. Mol Pharmacol, 2003. **63**(1): p. 9-18.
200. Kim, J., et al., *Functional antagonism of different G protein-coupled receptor kinases for beta-arrestin-mediated angiotensin II receptor signaling*. Proc Natl Acad Sci U S A, 2005. **102**(5): p. 1442-7.
201. Krasel, C., et al., *Beta-arrestin binding to the beta2-adrenergic receptor requires both receptor phosphorylation and receptor activation*. J Biol Chem, 2005. **280**(10): p. 9528-35.
202. Lefkowitz, R.J., *G protein-coupled receptors and receptor kinases: from molecular biology to potential therapeutic applications*. Nat Biotechnol, 1996. **14**(3): p. 283-6.
203. Mangmool, S., et al., *Clathrin required for phosphorylation and internalization of beta2-adrenergic receptor by G protein-coupled receptor kinase 2 (GRK2)*. J Biol Chem, 2006. **281**(42): p. 31940-9.
204. Shenoy, S.K., et al., *beta-arrestin-dependent, G protein-independent ERK1/2 activation by the beta2 adrenergic receptor*. J Biol Chem, 2006. **281**(2): p. 1261-73.
205. Simon, V., et al., *Endogenous G protein-coupled receptor kinase 6 triggers homologous beta-adrenergic receptor desensitization in primary uterine smooth muscle cells*. Endocrinology, 2003. **144**(7): p. 3058-66.
206. Hirsch, J.A., et al., *The 2.8 Å crystal structure of visual arrestin: a model for arrestin's regulation*. Cell, 1999. **97**(2): p. 257-69.
207. McDowell, J.H., J.P. Nawrocki, and P.A. Hargrave, *Phosphorylation sites in bovine rhodopsin*. Biochemistry, 1993. **32**(18): p. 4968-74.
208. Pulvermuller, A., et al., *Interactions of metarhodopsin II. Arrestin peptides compete with arrestin and transducin*. J Biol Chem, 2000. **275**(48): p. 37679-85.
209. Vishnivetskiy, S.A., et al., *Mapping the arrestin-receptor interface. Structural elements responsible for receptor specificity of arrestin proteins*. J Biol Chem, 2004. **279**(2): p. 1262-8.
210. Lefkowitz, R.J. and E.J. Whalen, *beta-arrestins: traffic cops of cell signaling*. Curr Opin Cell Biol, 2004. **16**(2): p. 162-8.

211. Zhang, M., et al., *Loss of betaarrestin1 and betaarrestin2 contributes to pulmonary hypoplasia and neonatal lethality in mice*. Dev Biol, 2010. **339**(2): p. 407-17.
212. Gurevich, V.V. and E.V. Gurevich, *The molecular acrobatics of arrestin activation*. Trends Pharmacol Sci, 2004. **25**(2): p. 105-11.
213. Conner, D.A., et al., *beta-Arrestin1 knockout mice appear normal but demonstrate altered cardiac responses to beta-adrenergic stimulation*. Circ Res, 1997. **81**(6): p. 1021-6.
214. Ahn, S., et al., *Desensitization, internalization, and signaling functions of beta-arrestins demonstrated by RNA interference*. Proc Natl Acad Sci U S A, 2003. **100**(4): p. 1740-4.
215. Lee, M.H., et al., *Role of beta-arrestin-mediated desensitization and signaling in the control of angiotensin AT1a receptor-stimulated transcription*. J Biol Chem, 2008. **283**(4): p. 2088-97.
216. Paing, M.M., et al., *beta -Arrestins regulate protease-activated receptor-1 desensitization but not internalization or Down-regulation*. J Biol Chem, 2002. **277**(2): p. 1292-300.
217. Kuo, F.T., T.L. Lu, and H.W. Fu, *Opposing effects of beta-arrestin1 and beta-arrestin2 on activation and degradation of Src induced by protease-activated receptor 1*. Cell Signal, 2006. **18**(11): p. 1914-23.
218. DeWire, S.M., et al., *Beta-arrestins and cell signaling*. Annu Rev Physiol, 2007. **69**: p. 483-510.
219. Oakley, R.H., et al., *Differential affinities of visual arrestin, beta arrestin1, and beta arrestin2 for G protein-coupled receptors delineate two major classes of receptors*. J Biol Chem, 2000. **275**(22): p. 17201-10.
220. Paing, M.M., et al., *Clathrin adaptor AP2 regulates thrombin receptor constitutive internalization and endothelial cell resensitization*. Mol Cell Biol, 2006. **26**(8): p. 3231-42.
221. Ge, L., et al., *A beta-arrestin-dependent scaffold is associated with prolonged MAPK activation in pseudopodia during protease-activated receptor-2-induced chemotaxis*. J Biol Chem, 2003. **278**(36): p. 34418-26.
222. Ge, L., et al., *Constitutive protease-activated receptor-2-mediated migration of MDA MB-231 breast cancer cells requires both beta-arrestin-1 and -2*. J Biol Chem, 2004. **279**(53): p. 55419-24.
223. Kumar, P., et al., *Differential effects of beta-arrestins on the internalization, desensitization and ERK1/2 activation downstream of protease activated receptor-2*. Am J Physiol Cell Physiol, 2007. **293**(1): p. C346-57.
224. Shenoy, S.K. and R.J. Lefkowitz, *beta-Arrestin-mediated receptor trafficking and signal transduction*. Trends Pharmacol Sci, 2011. **32**(9): p. 521-33.
225. Pierce, K.L. and R.J. Lefkowitz, *Classical and new roles of beta-arrestins in the regulation of G-protein-coupled receptors*. Nat Rev Neurosci, 2001. **2**(10): p. 727-33.
226. Shenoy, S.K. and R.J. Lefkowitz, *Multifaceted roles of beta-arrestins in the regulation of seven-membrane-spanning receptor trafficking and signalling*. Biochem J, 2003. **375**(Pt 3): p. 503-15.
227. Shenoy, S.K. and R.J. Lefkowitz, *Trafficking patterns of beta-arrestin and G protein-coupled receptors determined by the kinetics of beta-arrestin deubiquitination*. J Biol Chem, 2003. **278**(16): p. 14498-506.
228. Laporte, S.A., et al., *beta-Arrestin/AP-2 interaction in G protein-coupled receptor internalization: identification of a beta-arrestin binding site in beta 2-adaptin*. J Biol Chem, 2002. **277**(11): p. 9247-54.

229. Kim, Y.M. and J.L. Benovic, *Differential roles of arrestin-2 interaction with clathrin and adaptor protein 2 in G protein-coupled receptor trafficking*. J Biol Chem, 2002. **277**(34): p. 30760-8.
230. Luttrell, L.M., Y. Daaka, and R.J. Lefkowitz, *Regulation of tyrosine kinase cascades by G-protein-coupled receptors*. Curr Opin Cell Biol, 1999. **11**(2): p. 177-83.
231. Luttrell, L.M. and D. Gesty-Palmer, *Beyond desensitization: physiological relevance of arrestin-dependent signaling*. Pharmacol Rev, 2010. **62**(2): p. 305-30.
232. Ahn, S., et al., *Differential kinetic and spatial patterns of beta-arrestin and G protein-mediated ERK activation by the angiotensin II receptor*. J Biol Chem, 2004. **279**(34): p. 35518-25.
233. Tohgo, A., et al., *The stability of the G protein-coupled receptor-beta-arrestin interaction determines the mechanism and functional consequence of ERK activation*. J Biol Chem, 2003. **278**(8): p. 6258-67.
234. Lefkowitz, R.J. and S.K. Shenoy, *Transduction of receptor signals by beta-arrestins*. Science, 2005. **308**(5721): p. 512-7.
235. Lefkowitz, R.J., K. Rajagopal, and E.J. Whalen, *New roles for beta-arrestins in cell signaling: not just for seven-transmembrane receptors*. Mol Cell, 2006. **24**(5): p. 643-52.
236. Buchanan, F.G. and R.N. DuBois, *Emerging roles of beta-arrestins*. Cell Cycle, 2006. **5**(18): p. 2060-3.
237. Kang, J., et al., *A nuclear function of beta-arrestin1 in GPCR signaling: regulation of histone acetylation and gene transcription*. Cell, 2005. **123**(5): p. 833-47.
238. Wang, P., et al., *Subcellular localization of beta-arrestins is determined by their intact N domain and the nuclear export signal at the C terminus*. J Biol Chem, 2003. **278**(13): p. 11648-53.
239. McDonald, P.H., et al., *Beta-arrestin 2: a receptor-regulated MAPK scaffold for the activation of JNK3*. Science, 2000. **290**(5496): p. 1574-7.
240. Scott, M.G., et al., *Differential nucleocytoplasmic shuttling of beta-arrestins. Characterization of a leucine-rich nuclear export signal in beta-arrestin2*. J Biol Chem, 2002. **277**(40): p. 37693-701.
241. Shenoy, S.K., et al., *Beta-arrestin-dependent signaling and trafficking of 7-transmembrane receptors is reciprocally regulated by the deubiquitinase USP33 and the E3 ligase Mdm2*. Proc Natl Acad Sci U S A, 2009. **106**(16): p. 6650-5.
242. Wyatt, D., et al., *Small ubiquitin-like modifier modification of arrestin-3 regulates receptor trafficking*. J Biol Chem, 2011. **286**(5): p. 3884-93.
243. Lin, M.H., et al., *Clathrin-mediated endocytosis is required for ANE 30-100K-induced autophagy*. J Oral Pathol Med, 2017.
244. Lin, F.T., et al., *Clathrin-mediated endocytosis of the beta-adrenergic receptor is regulated by phosphorylation/dephosphorylation of beta-arrestin1*. J Biol Chem, 1997. **272**(49): p. 31051-7.
245. Lin, F.T., et al., *Feedback regulation of beta-arrestin1 function by extracellular signal-regulated kinases*. J Biol Chem, 1999. **274**(23): p. 15971-4.
246. Milano, S.K., et al., *Nonvisual arrestin oligomerization and cellular localization are regulated by inositol hexakisphosphate binding*. J Biol Chem, 2006. **281**(14): p. 9812-23.
247. Boularan, C., et al., *beta-arrestin 2 oligomerization controls the Mdm2-dependent inhibition of p53*. Proc Natl Acad Sci U S A, 2007. **104**(46): p. 18061-6.
248. Storez, H., et al., *Homo- and hetero-oligomerization of beta-arrestins in living cells*. J Biol Chem, 2005. **280**(48): p. 40210-5.

249. Kenakin, T., *Agonist-receptor efficacy. II. Agonist trafficking of receptor signals*. Trends Pharmacol Sci, 1995. **16**(7): p. 232-8.
250. McDonald, P.H. and R.J. Lefkowitz, *Beta-Arrestins: new roles in regulating heptahelical receptors' functions*. Cell Signal, 2001. **13**(10): p. 683-9.
251. Kenakin, T., *Ligand-selective receptor conformations revisited: the promise and the problem*. Trends Pharmacol Sci, 2003. **24**(7): p. 346-54.
252. Kahsai, A.W., et al., *Multiple ligand-specific conformations of the beta2-adrenergic receptor*. Nat Chem Biol, 2011. **7**(10): p. 692-700.
253. Violin, J.D. and R.J. Lefkowitz, *Beta-arrestin-biased ligands at seven-transmembrane receptors*. Trends Pharmacol Sci, 2007. **28**(8): p. 416-22.
254. Whalen, E.J., S. Rajagopal, and R.J. Lefkowitz, *Therapeutic potential of beta-arrestin- and G protein-biased agonists*. Trends Mol Med, 2011. **17**(3): p. 126-39.
255. Hupfeld, C.J. and J.M. Olefsky, *Regulation of receptor tyrosine kinase signaling by GRKs and beta-arrestins*. Annu Rev Physiol, 2007. **69**: p. 561-77.
256. Luttrell, L., et al., *A pertussis toxin-sensitive G-protein mediates some aspects of insulin action in BC3H-1 murine myocytes*. J Biol Chem, 1990. **265**(28): p. 16873-9.
257. Alderton, F., et al., *Tethering of the platelet-derived growth factor beta receptor to G-protein-coupled receptors. A novel platform for integrative signaling by these receptor classes in mammalian cells*. J Biol Chem, 2001. **276**(30): p. 28578-85.
258. Rakhit, S., S. Pyne, and N.J. Pyne, *Nerve growth factor stimulation of p42/p44 mitogen-activated protein kinase in PC12 cells: role of G(i/o), G protein-coupled receptor kinase 2, beta-arrestin I, and endocytic processing*. Mol Pharmacol, 2001. **60**(1): p. 63-70.
259. Xu, C.B., et al., *D-erythro-N,N-dimethylsphingosine inhibits bFGF-induced proliferation of cerebral, aortic and coronary smooth muscle cells*. Atherosclerosis, 2002. **164**(2): p. 237-43.
260. Kreuzer, J., et al., *Platelet-derived growth factor activates production of reactive oxygen species by NAD(P)H oxidase in smooth muscle cells through Gi1,2*. FASEB J, 2003. **17**(1): p. 38-40.
261. Zhang, B.H., V. Ho, and G.C. Farrell, *Specific involvement of G(alpha i2) with epidermal growth factor receptor signaling in rat hepatocytes, and the inhibitory effect of chronic ethanol*. Biochem Pharmacol, 2001. **61**(8): p. 1021-7.
262. Luttrell, L.M., et al., *G beta gamma subunits mediate mitogen-activated protein kinase activation by the tyrosine kinase insulin-like growth factor I receptor*. J Biol Chem, 1995. **270**(28): p. 16495-8.
263. Imamura, T., et al., *G alpha-q/11 protein plays a key role in insulin-induced glucose transport in 3T3-L1 adipocytes*. Mol Cell Biol, 1999. **19**(10): p. 6765-74.
264. Dalle, S., et al., *Insulin and insulin-like growth factor I receptors utilize different G protein signaling components*. J Biol Chem, 2001. **276**(19): p. 15688-95.
265. Delcourt, N., J. Bockaert, and P. Marin, *GPCR-jacking: from a new route in RTK signalling to a new concept in GPCR activation*. Trends Pharmacol Sci, 2007. **28**(12): p. 602-7.
266. Hallak, H., et al., *Association of heterotrimeric G(i) with the insulin-like growth factor-I receptor. Release of G(betagamma) subunits upon receptor activation*. J Biol Chem, 2000. **275**(4): p. 2255-8.
267. Gao, J., J. Li, and L. Ma, *Regulation of EGF-induced ERK/MAPK activation and EGFR internalization by G protein-coupled receptor kinase 2*. Acta Biochim Biophys Sin (Shanghai), 2005. **37**(8): p. 525-31.
268. Freedman, N.J., et al., *Phosphorylation of the platelet-derived growth factor receptor-beta and epidermal growth factor receptor by G protein-coupled receptor kinase-2*.

- Mechanisms for selectivity of desensitization.* J Biol Chem, 2002. **277**(50): p. 48261-9.
269. Zheng, H., et al., *Selective recruitment of G protein-coupled receptor kinases (GRKs) controls signaling of the insulin-like growth factor I receptor.* Proc Natl Acad Sci U S A, 2012. **109**(18): p. 7055-60.
  270. Lin, F.T., Y. Daaka, and R.J. Lefkowitz, *beta-arrestins regulate mitogenic signaling and clathrin-mediated endocytosis of the insulin-like growth factor I receptor.* J Biol Chem, 1998. **273**(48): p. 31640-3.
  271. Chow, J.C., G. Condorelli, and R.J. Smith, *Insulin-like growth factor-I receptor internalization regulates signaling via the Shc/mitogen-activated protein kinase pathway, but not the insulin receptor substrate-1 pathway.* J Biol Chem, 1998. **273**(8): p. 4672-80.
  272. Girnita, L., et al., *{beta}-Arrestin is crucial for ubiquitination and down-regulation of the insulin-like growth factor-I receptor by acting as adaptor for the MDM2 E3 ligase.* J Biol Chem, 2005. **280**(26): p. 24412-9.
  273. Girnita, L., A. Girnita, and O. Larsson, *Mdm2-dependent ubiquitination and degradation of the insulin-like growth factor I receptor.* Proc Natl Acad Sci U S A, 2003. **100**(14): p. 8247-52.
  274. Zhande, R., et al., *Molecular mechanism of insulin-induced degradation of insulin receptor substrate 1.* Mol Cell Biol, 2002. **22**(4): p. 1016-26.
  275. Rui, L., et al., *SOCS-1 and SOCS-3 block insulin signaling by ubiquitin-mediated degradation of IRS1 and IRS2.* J Biol Chem, 2002. **277**(44): p. 42394-8.
  276. Usui, I., et al., *beta-arrestin-1 competitively inhibits insulin-induced ubiquitination and degradation of insulin receptor substrate 1.* Mol Cell Biol, 2004. **24**(20): p. 8929-37.
  277. Dalle, S., et al., *Insulin induces heterologous desensitization of G-protein-coupled receptor and insulin-like growth factor I signaling by downregulating beta-arrestin-1.* Mol Cell Biol, 2002. **22**(17): p. 6272-85.
  278. Lee, A.V., et al., *Insulin-like growth factor I-induced degradation of insulin receptor substrate 1 is mediated by the 26S proteasome and blocked by phosphatidylinositol 3'-kinase inhibition.* Mol Cell Biol, 2000. **20**(5): p. 1489-96.
  279. Kim, J., et al., *Regulation of epidermal growth factor receptor internalization by G protein-coupled receptors.* Biochemistry, 2003. **42**(10): p. 2887-94.
  280. Hobson, J.P., et al., *Role of the sphingosine-1-phosphate receptor EDG-1 in PDGF-induced cell motility.* Science, 2001. **291**(5509): p. 1800-3.
  281. Waters, C., S. Pyne, and N.J. Pyne, *The role of G-protein coupled receptors and associated proteins in receptor tyrosine kinase signal transduction.* Semin Cell Dev Biol, 2004. **15**(3): p. 309-23.
  282. Girnita, L., et al., *Beta-arrestin and Mdm2 mediate IGF-1 receptor-stimulated ERK activation and cell cycle progression.* J Biol Chem, 2007. **282**(15): p. 11329-38.
  283. Hunton, D.L., et al., *Beta-arrestin 2-dependent angiotensin II type 1A receptor-mediated pathway of chemotaxis.* Mol Pharmacol, 2005. **67**(4): p. 1229-36.
  284. Buchanan, F.G., et al., *Role of beta-arrestin 1 in the metastatic progression of colorectal cancer.* Proc Natl Acad Sci U S A, 2006. **103**(5): p. 1492-7.
  285. Luan, B., et al., *Deficiency of a beta-arrestin-2 signal complex contributes to insulin resistance.* Nature, 2009. **457**(7233): p. 1146-9.
  286. Povsic, T.J., T.A. Kohout, and R.J. Lefkowitz, *Beta-arrestin1 mediates insulin-like growth factor 1 (IGF-1) activation of phosphatidylinositol 3-kinase (PI3K) and anti-apoptosis.* J Biol Chem, 2003. **278**(51): p. 51334-9.



287. De Smet, F., A. Christopoulos, and P. Carmeliet, *Allosteric targeting of receptor tyrosine kinases*. Nat Biotechnol, 2014. **32**(11): p. 1113-20.
288. Karaman, M.W., et al., *A quantitative analysis of kinase inhibitor selectivity*. Nat Biotechnol, 2008. **26**(1): p. 127-32.
289. Crudden, C., et al., *The dichotomy of the Insulin-like growth factor 1 receptor: RTK and GPCR: friend or foe for cancer treatment?* Growth Horm IGF Res, 2014.
290. Laine, E., C. Auclair, and L. Tchertanov, *Allosteric communication across the native and mutated KIT receptor tyrosine kinase*. PLoS Comput Biol, 2012. **8**(8): p. e1002661.
291. Hyde, C.A., et al., *Targeting extracellular domains D4 and D7 of vascular endothelial growth factor receptor 2 reveals allosteric receptor regulatory sites*. Mol Cell Biol, 2012. **32**(19): p. 3802-13.
292. Cazorla, M., et al., *Cyclotraxin-B, the first highly potent and selective TrkB inhibitor, has anxiolytic properties in mice*. PLoS One, 2010. **5**(3): p. e9777.
293. Udugamasooriya, D.G., et al., *A peptoid "antibody surrogate" that antagonizes VEGF receptor 2 activity*. J Am Chem Soc, 2008. **130**(17): p. 5744-52.
294. Tvorogov, D., et al., *Effective suppression of vascular network formation by combination of antibodies blocking VEGFR ligand binding and receptor dimerization*. Cancer Cell, 2010. **18**(6): p. 630-40.
295. Leahy, D.J., *A molecular view of anti-ErbB monoclonal antibody therapy*. Cancer Cell, 2008. **13**(4): p. 291-3.
296. Fleishman, S.J., J. Schlessinger, and N. Ben-Tal, *A putative molecular-activation switch in the transmembrane domain of erbB2*. Proc Natl Acad Sci U S A, 2002. **99**(25): p. 15937-40.
297. Weisheit, S., et al., *PKCepsilon acts as negative allosteric modulator of EGF receptor signalling*. Cell Signal, 2011. **23**(2): p. 436-48.
298. Weisheit, S. and C. Liebmann, *Allosteric modulation by protein kinase Cepsilon leads to modified responses of EGF receptor towards tyrosine kinase inhibitors*. Cell Signal, 2012. **24**(2): p. 422-34.
299. Grunewald, F.S., et al., *Structure-function analysis of VEGF receptor activation and the role of coreceptors in angiogenic signaling*. Biochim Biophys Acta, 2010. **1804**(3): p. 567-80.
300. Landgraf, K.E., et al., *Allosteric peptide activators of pro-hepatocyte growth factor stimulate Met signaling*. J Biol Chem, 2010. **285**(51): p. 40362-72.
301. Corbin, J.A., et al., *Improved glucose metabolism in vitro and in vivo by an allosteric monoclonal antibody that increases insulin receptor binding affinity*. PLoS One, 2014. **9**(2): p. e88684.
302. Bono, F., et al., *Inhibition of tumor angiogenesis and growth by a small-molecule multi-FGF receptor blocker with allosteric properties*. Cancer Cell, 2013. **23**(4): p. 477-88.
303. Herbert, C., et al., *Molecular mechanism of SSR128129E, an extracellularly acting, small-molecule, allosteric inhibitor of FGF receptor signaling*. Cancer Cell, 2013. **23**(4): p. 489-501.
304. Jang, S.W., et al., *Gambogic amide, a selective agonist for TrkA receptor that possesses robust neurotrophic activity, prevents neuronal cell death*. Proc Natl Acad Sci U S A, 2007. **104**(41): p. 16329-34.
305. Vasilcanu, R., et al., *Insulin-like growth factor type-I receptor-dependent phosphorylation of extracellular signal-regulated kinase 1/2 but not Akt (protein kinase B) can be induced by picropodophyllin*. Mol Pharmacol, 2008. **73**(3): p. 930-9.

306. Girnita, A., et al., *Identification of the cathelicidin peptide LL-37 as agonist for the type I insulin-like growth factor receptor*. *Oncogene*, 2012. **31**(3): p. 352-65.
307. Zheng, H., et al., *beta-Arrestin-biased agonism as the central mechanism of action for insulin-like growth factor 1 receptor-targeting antibodies in Ewing's sarcoma*. *Proc Natl Acad Sci U S A*, 2012. **109**(50): p. 20620-5.
308. Dasgupta, P., et al., *ARRB1-mediated regulation of E2F target genes in nicotine-induced growth of lung tumors*. *J Natl Cancer Inst*, 2011. **103**(4): p. 317-33.
309. Revankar, C.M., et al., *Arrestins block G protein-coupled receptor-mediated apoptosis*. *J Biol Chem*, 2004. **279**(23): p. 24578-84.
310. Fereshteh, M., et al., *beta-Arrestin2 mediates the initiation and progression of myeloid leukemia*. *Proc Natl Acad Sci U S A*, 2012. **109**(31): p. 12532-7.
311. Rosano, L., et al., *Beta-arrestin links endothelin A receptor to beta-catenin signaling to induce ovarian cancer cell invasion and metastasis*. *Proc Natl Acad Sci U S A*, 2009. **106**(8): p. 2806-11.
312. Wu, Z., et al., *[The clinical significance of beta-arrestin 2 expression in the serum of non-small cell lung cancer patients]*. *Zhongguo Fei Ai Za Zhi*, 2011. **14**(6): p. 497-501.
313. Bonnans, C., et al., *Essential requirement for beta-arrestin2 in mouse intestinal tumors with elevated Wnt signaling*. *Proc Natl Acad Sci U S A*, 2012. **109**(8): p. 3047-52.
314. Sobolesky, P.M. and O. Moussa, *The role of beta-arrestins in cancer*. *Prog Mol Biol Transl Sci*, 2013. **118**: p. 395-411.
315. Beaulieu, J.M. and M.G. Caron, *Beta-arrestin goes nuclear*. *Cell*, 2005. **123**(5): p. 755-7.
316. Chen, W., et al., *beta-Arrestin1 modulates lymphoid enhancer factor transcriptional activity through interaction with phosphorylated dishevelled proteins*. *Proc Natl Acad Sci U S A*, 2001. **98**(26): p. 14889-94.
317. Zou, L., et al., *Rapid xenograft tumor progression in beta-arrestin1 transgenic mice due to enhanced tumor angiogenesis*. *FASEB J*, 2008. **22**(2): p. 355-64.
318. DeFea, K.A., et al., *The proliferative and antiapoptotic effects of substance P are facilitated by formation of a beta -arrestin-dependent scaffolding complex*. *Proc Natl Acad Sci U S A*, 2000. **97**(20): p. 11086-91.
319. Wagener, B.M., et al., *Adaptor protein-2 interaction with arrestin regulates GPCR recycling and apoptosis*. *Traffic*, 2009. **10**(9): p. 1286-300.
320. Vines, C.M., et al., *N-formyl peptide receptors internalize but do not recycle in the absence of arrestins*. *J Biol Chem*, 2003. **278**(43): p. 41581-4.
321. Ahn, S., et al., *{beta}-Arrestin-2 Mediates Anti-apoptotic Signaling through Regulation of BAD Phosphorylation*. *J Biol Chem*, 2009. **284**(13): p. 8855-65.
322. Li, H., et al., *beta-arrestin 2 regulates Toll-like receptor 4-mediated apoptotic signalling through glycogen synthase kinase-3beta*. *Immunology*, 2010. **130**(4): p. 556-63.
323. Sun, X., et al., *Beta-arrestin 2 modulates resveratrol-induced apoptosis and regulation of Akt/GSK3ss pathways*. *Biochim Biophys Acta*, 2010. **1800**(9): p. 912-8.
324. Fong, A.M., et al., *Defective lymphocyte chemotaxis in beta-arrestin2- and GRK6-deficient mice*. *Proc Natl Acad Sci U S A*, 2002. **99**(11): p. 7478-83.
325. Sun, Y., et al., *Beta-arrestin2 is critically involved in CXCR4-mediated chemotaxis, and this is mediated by its enhancement of p38 MAPK activation*. *J Biol Chem*, 2002. **277**(51): p. 49212-9.
326. Luan, B., et al., *Beta-arrestin2 functions as a phosphorylation-regulated suppressor of UV-induced NF-kappaB activation*. *EMBO J*, 2005. **24**(24): p. 4237-46.

327. Wang, P., et al., *Beta-arrestin 2 functions as a G-protein-coupled receptor-activated regulator of oncoprotein Mdm2*. J Biol Chem, 2003. **278**(8): p. 6363-70.
328. Moussa, O., et al., *Novel role of thromboxane receptors beta isoform in bladder cancer pathogenesis*. Cancer Res, 2008. **68**(11): p. 4097-104.
329. Laroche, G., et al., *Involvement of actin in agonist-induced endocytosis of the G protein-coupled receptor for thromboxane A2: overcoming of actin disruption by arrestin-3 but not arrestin-2*. J Biol Chem, 2005. **280**(24): p. 23215-24.
330. Liu, H., et al., *Elevated beta-arrestin1 expression correlated with risk stratification in acute lymphoblastic leukemia*. Int J Hematol, 2011. **93**(4): p. 494-501.
331. Duan, X., et al., *beta-arrestin1 promotes epithelial-mesenchymal transition via modulating GSK-3beta/beta-catenin pathway in prostate cancer cells*. Biochem Biophys Res Commun, 2016. **479**(2): p. 204-210.
332. Lakshmikanthan, V., et al., *Identification of betaArrestin2 as a corepressor of androgen receptor signaling in prostate cancer*. Proc Natl Acad Sci U S A, 2009. **106**(23): p. 9379-84.
333. Turley, R.S., et al., *The type III transforming growth factor-beta receptor as a novel tumor suppressor gene in prostate cancer*. Cancer Res, 2007. **67**(3): p. 1090-8.
334. Mythreye, K. and G.C. Blobe, *The type III TGF-beta receptor regulates epithelial and cancer cell migration through beta-arrestin2-mediated activation of Cdc42*. Proc Natl Acad Sci U S A, 2009. **106**(20): p. 8221-6.
335. Zhang, P., et al., *beta-arrestin2 mediates beta-2 adrenergic receptor signaling inducing prostate cancer cell progression*. Oncol Rep, 2011. **26**(6): p. 1471-7.
336. Shen, X., et al., *Chemokine receptor CXCR4 enhances proliferation in pancreatic cancer cells through AKT and ERK dependent pathways*. Pancreas, 2010. **39**(1): p. 81-7.
337. Heinrich, E.L., et al., *Chemokine CXCL12 activates dual CXCR4 and CXCR7-mediated signaling pathways in pancreatic cancer cells*. J Transl Med, 2012. **10**: p. 68.
338. Chen, W., et al., *Activity-dependent internalization of smoothened mediated by beta-arrestin 2 and GRK2*. Science, 2004. **306**(5705): p. 2257-60.
339. Thayer, S.P., et al., *Hedgehog is an early and late mediator of pancreatic cancer tumorigenesis*. Nature, 2003. **425**(6960): p. 851-6.
340. Mandell, J.W., et al., *Dephosphorylation of beta-arrestin 1 in glioblastomas*. J Neuropathol Exp Neurol, 2009. **68**(5): p. 535-41.
341. Decaillot, F.M., et al., *CXCR7/CXCR4 heterodimer constitutively recruits beta-arrestin to enhance cell migration*. J Biol Chem, 2011. **286**(37): p. 32188-97.
342. Kim, J.I., et al., *Prostaglandin E2 promotes lung cancer cell migration via EP4-betaArrestin1-c-Src signalsome*. Mol Cancer Res, 2010. **8**(4): p. 569-77.
343. Dasgupta, P., et al., *Nicotine induces cell proliferation by beta-arrestin-mediated activation of Src and Rb-Raf-1 pathways*. J Clin Invest, 2006. **116**(8): p. 2208-2217.
344. Li, T.T., et al., *Beta-arrestin/Ral signaling regulates lysophosphatidic acid-mediated migration and invasion of human breast tumor cells*. Mol Cancer Res, 2009. **7**(7): p. 1064-77.
345. Shenoy, S.K., et al., *beta-arrestin1 mediates metastatic growth of breast cancer cells by facilitating HIF-1-dependent VEGF expression*. Oncogene, 2012. **31**(3): p. 282-92.
346. Michal, A.M., et al., *Differential expression of arrestins is a predictor of breast cancer progression and survival*. Breast Cancer Res Treat, 2011. **130**(3): p. 791-807.
347. Goertzen, C.G., et al., *KISS1R signaling promotes invadopodia formation in human breast cancer cell via beta-arrestin2/ERK*. Cell Signal, 2016. **28**(3): p. 165-76.

348. Ray, P., et al., *CXCL12-gamma in primary tumors drives breast cancer metastasis*. *Oncogene*, 2015. **34**(16): p. 2043-51.
349. You, H.J., T. How, and G.C. Blobe, *The type III transforming growth factor-beta receptor negatively regulates nuclear factor kappa B signaling through its interaction with beta-arrestin2*. *Carcinogenesis*, 2009. **30**(8): p. 1281-7.
350. Bedi, A. and B. Mookerjee, *Biological significance and molecular mechanisms of p53-induced apoptosis*. *Apoptosis*, 1998. **3**(4): p. 237-44.
351. Berns, A., *Cancer genetics. Is p53 the only real tumor suppressor gene?* *Curr Biol*, 1994. **4**(2): p. 137-9.
352. Bennett, M.R., *Mechanisms of p53-induced apoptosis*. *Biochem Pharmacol*, 1999. **58**(7): p. 1089-95.
353. Oren, M., *p53: the ultimate tumor suppressor gene?* *FASEB J*, 1992. **6**(13): p. 3169-76.
354. Vousden, K.H. and C. Prives, *Blinded by the Light: The Growing Complexity of p53*. *Cell*, 2009. **137**(3): p. 413-31.
355. Gorgoulis, V.G., et al., *Activation of the DNA damage checkpoint and genomic instability in human precancerous lesions*. *Nature*, 2005. **434**(7035): p. 907-13.
356. Bartkova, J., et al., *DNA damage response as a candidate anti-cancer barrier in early human tumorigenesis*. *Nature*, 2005. **434**(7035): p. 864-70.
357. Levine, A.J., *p53, the cellular gatekeeper for growth and division*. *Cell*, 1997. **88**(3): p. 323-31.
358. Soussi, T. and K.G. Wiman, *Shaping genetic alterations in human cancer: the p53 mutation paradigm*. *Cancer Cell*, 2007. **12**(4): p. 303-12.
359. Soussi, T. and K.G. Wiman, *TP53: an oncogene in disguise*. *Cell Death Differ*, 2015. **22**(8): p. 1239-49.
360. Leach, F.S., et al., *p53 Mutation and MDM2 amplification in human soft tissue sarcomas*. *Cancer Res*, 1993. **53**(10 Suppl): p. 2231-4.
361. Bond, G.L., et al., *A single nucleotide polymorphism in the MDM2 promoter attenuates the p53 tumor suppressor pathway and accelerates tumor formation in humans*. *Cell*, 2004. **119**(5): p. 591-602.
362. Molchadsky, A., et al., *p53 is required for brown adipogenic differentiation and has a protective role against diet-induced obesity*. *Cell Death Differ*, 2013. **20**(5): p. 774-83.
363. Schwartzberg-Bar-Yoseph, F., M. Armoni, and E. Karnieli, *The tumor suppressor p53 down-regulates glucose transporters GLUT1 and GLUT4 gene expression*. *Cancer Res*, 2004. **64**(7): p. 2627-33.
364. Werner, H. and S. Maor, *The insulin-like growth factor-I receptor gene: a downstream target for oncogene and tumor suppressor action*. *Trends Endocrinol Metab*, 2006. **17**(6): p. 236-42.
365. Ohlsson, C., et al., *p53 regulates insulin-like growth factor-I (IGF-I) receptor expression and IGF-I-induced tyrosine phosphorylation in an osteosarcoma cell line: interaction between p53 and Sp1*. *Endocrinology*, 1998. **139**(3): p. 1101-7.
366. Werner, H., et al., *Wild-type and mutant p53 differentially regulate transcription of the insulin-like growth factor I receptor gene*. *Proc Natl Acad Sci U S A*, 1996. **93**(16): p. 8318-23.
367. Zhang, L., et al., *Regulation of insulin-like growth factor II P3 promotor by p53: a potential mechanism for tumorigenesis*. *Cancer Res*, 1996. **56**(6): p. 1367-73.
368. Buckbinder, L., et al., *Induction of the growth inhibitor IGF-binding protein 3 by p53*. *Nature*, 1995. **377**(6550): p. 646-9.

369. Grimberg, A., et al., *Insulin-like growth factor factor binding protein-2 is a novel mediator of p53 inhibition of insulin-like growth factor signaling*. *Cancer Biol Ther*, 2006. **5**(10): p. 1408-14.
370. Nahor, I., et al., *The p53-family members p63 and p73 inhibit insulin-like growth factor-I receptor gene expression in colon cancer cells*. *Growth Horm IGF Res*, 2005. **15**(6): p. 388-96.
371. Maor, S.B., et al., *BRCA1 suppresses insulin-like growth factor-I receptor promoter activity: potential interaction between BRCA1 and Sp1*. *Mol Genet Metab*, 2000. **69**(2): p. 130-6.
372. Abramovitch, S. and H. Werner, *Functional and physical interactions between BRCA1 and p53 in transcriptional regulation of the IGF-IR gene*. *Horm Metab Res*, 2003. **35**(11-12): p. 758-62.
373. Yuen, J.S., et al., *The VHL tumor suppressor inhibits expression of the IGF1R and its loss induces IGF1R upregulation in human clear cell renal carcinoma*. *Oncogene*, 2007. **26**(45): p. 6499-508.
374. Werner, H., et al., *Inhibition of cellular proliferation by the Wilms' tumor suppressor WT1 is associated with suppression of insulin-like growth factor I receptor gene expression*. *Mol Cell Biol*, 1995. **15**(7): p. 3516-22.
375. Werner, H., et al., *Insulin-like Growth Factor I Signaling Axis Meets p53 Genome Protection Pathways*. *Front Oncol*, 2016. **6**: p. 159.
376. LeRoith, D. and S. Yakar, *Mechanisms of disease: metabolic effects of growth hormone and insulin-like growth factor I*. *Nat Clin Pract Endocrinol Metab*, 2007. **3**(3): p. 302-10.
377. Liao, Y., et al., *Up-regulation of insulin-like growth factor axis components in human primary prostate cancer correlates with tumor grade*. *Hum Pathol*, 2005. **36**(11): p. 1186-96.
378. Goldstein, I., et al., *p53 promotes the expression of gluconeogenesis-related genes and enhances hepatic glucose production*. *Cancer Metab*, 2013. **1**(1): p. 9.
379. Smith, T.M., et al., *IGF-I induces SREBP-1 expression and lipogenesis in SEB-I sebocytes via activation of the phosphoinositide 3-kinase/Akt pathway*. *J Invest Dermatol*, 2008. **128**(5): p. 1286-93.
380. Yahagi, N., et al., *p53 Activation in adipocytes of obese mice*. *J Biol Chem*, 2003. **278**(28): p. 25395-400.
381. Lakin, N.D. and S.P. Jackson, *Regulation of p53 in response to DNA damage*. *Oncogene*, 1999. **18**(53): p. 7644-55.
382. Brunet, A., et al., *Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor*. *Cell*, 1999. **96**(6): p. 857-68.
383. Zhou, B.P., et al., *HER-2/neu induces p53 ubiquitination via Akt-mediated MDM2 phosphorylation*. *Nat Cell Biol*, 2001. **3**(11): p. 973-82.
384. Heron-Milhavet, L. and D. LeRoith, *Insulin-like growth factor I induces MDM2-dependent degradation of p53 via the p38 MAPK pathway in response to DNA damage*. *J Biol Chem*, 2002. **277**(18): p. 15600-6.
385. Mayo, L.D. and D.B. Donner, *A phosphatidylinositol 3-kinase/Akt pathway promotes translocation of Mdm2 from the cytoplasm to the nucleus*. *Proc Natl Acad Sci U S A*, 2001. **98**(20): p. 11598-603.
386. Ganguli, G. and B. Wasylyk, *p53-independent functions of MDM2*. *Mol Cancer Res*, 2003. **1**(14): p. 1027-35.
387. Feng, Z., et al., *The regulation of AMPK beta1, TSC2, and PTEN expression by p53: stress, cell and tissue specificity, and the role of these gene products in modulating the IGF-I-AKT-mTOR pathways*. *Cancer Res*, 2007. **67**(7): p. 3043-53.

388. Leri, A., et al., *Insulin-like growth factor-1 induces Mdm2 and down-regulates p53, attenuating the myocyte renin-angiotensin system and stretch-mediated apoptosis*. Am J Pathol, 1999. **154**(2): p. 567-80.
389. Heron-Milhavet, L., et al., *Insulin-like growth factor-I (IGF-I) receptor activation rescues UV-damaged cells through a p38 signaling pathway. Potential role of the IGF-I receptor in DNA repair*. J Biol Chem, 2001. **276**(21): p. 18185-92.
390. Takahashi, K. and K. Suzuki, *Association of insulin-like growth-factor-I-induced DNA synthesis with phosphorylation and nuclear exclusion of p53 in human breast cancer MCF-7 cells*. Int J Cancer, 1993. **55**(3): p. 453-8.
391. Marchetti, A., et al., *p53 can inhibit cell proliferation through caspase-mediated cleavage of ERK2/MAPK*. Cell Death Differ, 2004. **11**(6): p. 596-607.
392. Hara, M.R., et al., *A stress response pathway regulates DNA damage through beta2-adrenoreceptors and beta-arrestin-1*. Nature, 2011. **477**(7364): p. 349-53.
393. Castresana, J.S., et al., *Lack of allelic deletion and point mutation as mechanisms of p53 activation in human malignant melanoma*. Int J Cancer, 1993. **55**(4): p. 562-5.
394. Albino, A.P., et al., *Mutation and expression of the p53 gene in human malignant melanoma*. Melanoma Res, 1994. **4**(1): p. 35-45.
395. Hocker, T. and H. Tsao, *Ultraviolet radiation and melanoma: a systematic review and analysis of reported sequence variants*. Hum Mutat, 2007. **28**(6): p. 578-88.
396. Fang, S., et al., *Mdm2 is a RING finger-dependent ubiquitin protein ligase for itself and p53*. J Biol Chem, 2000. **275**(12): p. 8945-51.
397. Lee, J.T., P. Brafford, and M. Herlyn, *Unraveling the mysteries of IGF-1 signaling in melanoma*. J Invest Dermatol, 2008. **128**(6): p. 1358-60.
398. Capoluongo, E., *Insulin-like growth factor system and sporadic malignant melanoma*. Am J Pathol, 2011. **178**(1): p. 26-31.
399. Neudauer, C.L. and J.B. McCarthy, *Insulin-like growth factor I-stimulated melanoma cell migration requires phosphoinositide 3-kinase but not extracellular-regulated kinase activation*. Exp Cell Res, 2003. **286**(1): p. 128-37.
400. Shtivelman, E., et al., *Pathways and therapeutic targets in melanoma*. Oncotarget, 2014. **5**(7): p. 1701-52.
401. Villanueva, J., A. Vultur, and M. Herlyn, *Resistance to BRAF inhibitors: unraveling mechanisms and future treatment options*. Cancer Res, 2011. **71**(23): p. 7137-40.
402. Welsh, S.J., et al., *Resistance to combination BRAF and MEK inhibition in metastatic melanoma: Where to next?* Eur J Cancer, 2016. **62**: p. 76-85.
403. Spagnolo, F., P. Ghiorzo, and P. Queirolo, *Overcoming resistance to BRAF inhibition in BRAF-mutated metastatic melanoma*. Oncotarget, 2014. **5**(21): p. 10206-21.
404. Miller, M.A., et al., *Reduced Proteolytic Shedding of Receptor Tyrosine Kinases Is a Post-Translational Mechanism of Kinase Inhibitor Resistance*. Cancer Discov, 2016. **6**(4): p. 382-99.
405. Villanueva, J., et al., *Acquired resistance to BRAF inhibitors mediated by a RAF kinase switch in melanoma can be overcome by cotargeting MEK and IGF-1R/PI3K*. Cancer Cell, 2010. **18**(6): p. 683-95.
406. Herkert, B., et al., *Maximizing the Efficacy of MAPK-Targeted Treatment in PTENLOF/BRAFMUT Melanoma through PI3K and IGF1R Inhibition*. Cancer Res, 2016. **76**(2): p. 390-402.
407. Vasilcanu, R., et al., *Picropodophyllin induces downregulation of the insulin-like growth factor 1 receptor: potential mechanistic involvement of Mdm2 and beta-arrestin1*. Oncogene, 2008. **27**(11): p. 1629-38.
408. Baserga, R., *The insulin-like growth factor-I receptor as a target for cancer therapy*. Expert Opin Ther Targets, 2005. **9**(4): p. 753-68.

409. Yeh, A.H., E.A. Bohula, and V.M. Macaulay, *Human melanoma cells expressing V600E B-RAF are susceptible to IGF1R targeting by small interfering RNAs*. *Oncogene*, 2006. **25**(50): p. 6574-81.
410. Durfort, T., et al., *Small interfering RNA targeted to IGF-1R delays tumor growth and induces proinflammatory cytokines in a mouse breast cancer model*. *PLoS One*, 2012. **7**(1): p. e29213.
411. Worrall, C., et al., *Unbalancing p53/Mdm2/IGF-1R axis by Mdm2 activation restrains the IGF-1-dependent invasive phenotype of skin melanoma*. *Oncogene*, 2017.
412. Suleymanova, N., et al., *Functional antagonism of beta-arrestin isoforms balance IGF-1R expression and signalling with distinct cancer-related biological outcomes*. *Oncogene*, 2017.
413. Kondo, N., et al., *DNA damage induced by alkylating agents and repair pathways*. *J Nucleic Acids*, 2010. **2010**: p. 543531.
414. Sterne-Marr, R., et al., *Polypeptide variants of beta-arrestin and arrestin3*. *J Biol Chem*, 1993. **268**(21): p. 15640-8.
415. Gurevich, V.V. and E.V. Gurevich, *Structural determinants of arrestin functions*. *Prog Mol Biol Transl Sci*, 2013. **118**: p. 57-92.
416. Hara, M.R., et al., *Pharmacological blockade of a beta(2)AR-beta-arrestin-1 signaling cascade prevents the accumulation of DNA damage in a behavioral stress model*. *Cell Cycle*, 2013. **12**(2): p. 219-24.